

Caecal Gurgling - A new sign in the diagnosis of acute appendicitis and its correlation with intra operative findings



Susavan Das¹, Shib Shankar Kuri², Sayak Roy³, Kanchan Kundu⁴

^{1,3,4}Senior Resident, ²Associate Professor, Department of General Surgery, Bankura Sammilani Medical College, Bankura, West Bengal, India

Submission: 08-01-2022

Revision: 05-03-2022

Publication: 01-04-2022

ABSTRACT

Background: Acute appendicitis is one of the most common surgical emergencies, with a lifetime prevalence rate of approximately one in seven. The incidence is 1.5–1.9/1,000 in the male and female population and is approximately 1.4 times greater in men than in women. The diagnosis of acute appendicitis is based purely on clinical history and examination combined with laboratory investigations such as elevated white cell count. **Aims and Objectives:** To examine clinical and laboratory findings of patients with right iliac fossa pain attending general surgery outdoor, indoor, and emergency of BSMCH and the relation between caecal gurgling with acute appendicitis and its correlation with intraoperative findings. **Materials and Methods:** This study was a hospital-based cross-sectional analytical study. It was conducted in rural-based tertiary care hospital and medical college with a time frame of about 1½ years from acceptance of synopsis. The period of study was from January 2019 to May 2020, in the Department of general surgery, Bankura Sammilani Medical College and Hospital, Bankura. **Results:** In Acute Appendicitis, 188(68.1%) patients were APL, and 88(31.9%) patients were BPL. In Normal Appendix, 85(68.5%) patients were APL and 39(31.5%) patient were BPL. Association of SES versus The final diagnosis was not statistically significant (P=0.9315). **Conclusion:** Diagnosis of acute appendicitis is mainly clinical. Keeping in view the results of the present study, we conclude that caecal gurgling is an important positive clinical finding in the diagnosis of acute appendicitis. Incorporating caecal gurgling into Modified Alvarado Scoring System, especially in the patients with a MAS of 5–6 where there is a dilemma whether to operate the patient or not, it aids in our decision-making and thus helps to reduce the rate of negative appendectomies.

Key words: Acute appendicitis; Caecal gurgling; Intra operative findings

INTRODUCTION

Acute appendicitis is one of the most common surgical emergencies, with a lifetime prevalence rate of approximately one in seven.¹ The incidence is 1.5–1.9/1,000 in the male and female population and is approximately 1.4 times greater in men than in women.²

The diagnosis of acute appendicitis is based purely on clinical history and examination combined with laboratory investigations such as elevated white cell count. Despite being a common problem, acute

appendicitis remains a difficult diagnosis to establish, particularly among the young, the elderly, and females of reproductive age, where a host of other genitourinary and gynecological inflammatory conditions can present with signs and symptoms that are similar to those of acute appendicitis.³

A delay in performing an appendectomy in order to improve its diagnostic accuracy increases the risk of appendicular perforation and sepsis, which in turn increases morbidity and mortality. The opposite is also true, where with reduced diagnostic accuracy, the negative or

Access this article online

Website:

<http://nepjol.info/index.php/AJMS>

DOI:10.3126/ajms.v13i4.42156

E-ISSN: 2091-0576

P-ISSN: 2467-9100

Copyright (c) 2022 Asian Journal of Medical Sciences



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

Address for Correspondence:

Dr. Kanchan Kundu, Paschim Pranabananda Pally, Senior Resident, Department of General Surgery, Bankura Sammilani Medical College, Bankura – 722 102, West Bengal, India. **Mobile:** 9153832572. **E-mail:** kanchankundubsmc@gmail.com

unnecessary appendectomy rate is increased, and this is generally reported to be approximately 20–40%.

Diagnostic accuracy can be further improved through the use of ultrasonography or computed tomography imaging.⁴ However, these modalities are costly and may not be easily available all the time. Making arrangements for these diagnostic modalities may lead to further delays in diagnosis and surgery.

Symptoms of ileus caused by the direct invasion of the intestinal walls by the infective microbes or their toxic products, or there may be true obstruction may be due to kinks or strangulation produced by the inflammatory exudates. The characteristic picture of ileus then develops and clinically a palpable gurgling is present.³

Cecal gurgling, a relatively new sign, which can aid Modified ALVARADO scoring system in individuals with a score of 5–6, that is, in the observational group, in predicting acute appendicitis and whether to operate it or not. The purpose of this study is to validate the sign and its correlation with intraoperative findings in our set up.

Aims and objectives

General objective

To find out predictive validity of caecal gurgling in patients with Modified Alvarado score of 5–6 for diagnosis of acute appendicitis.

Specific objective

To examine clinical and laboratory findings of patients with right iliac fossa pain attending general surgery outdoor, indoor, and emergency of BSMCH and the relation between caecal gurgling with acute appendicitis and its correlation with intraoperative findings.

MATERIALS AND METHODS

The current study was a hospital-based cross-sectional analytical study, conducted in rural-based tertiary care hospital and medical college with a time frame of about 1½ years from acceptance of synopsis.

Study population

All patients attending surgery OPD and emergency with pain in right iliac fossa with a Modified Alvarado score of <7 and operated of Bankura Sammilani Medical College and Hospital, with inclusion criteria and exclusion criteria applied properly. According to previous census, the number of patients should be around 400.

Sample size

All patients presenting with complaints of right iliac fossa pain and other symptoms and signs suggestive of acute

appendicitis have been picked up for 5 days a week. The 5 days of each week have been chosen randomly.

Study design: Census method

Case, control - required or not:-

Case: Patients with Modified Alvarado score of <7 with cecal gurgling and operated

Control: Patients with Modified Alvarado score of <7 without cecal gurgling but operated

Inclusion criteria

1. All patients presenting with right iliac fossa pain in the age group of 13–60 years.

Exclusion criteria

1. History of RIF pain for >48 h.
2. Patients with a palpable lump at RIF.
3. Patients with features of appendicular abscess and perforation
4. Those who have been admitted by other specialties for other complaints but subsequently developed RIF pain.
5. Patients presenting with proven malignancy.
6. Patients with a history of trauma over the abdomen.
7. Patients with pregnancy and other comorbid conditions.

RESULTS

In our study among 400 patients, 306(76.5%) patients had Cecal gurgling and 94 patients were without cecal gurgling.

347(86.8%) patients had Intraoperative findings of Inflamed Appendix and 53(13.3%) patients were with Normal Appendix.

325(81.3%) patients had Inflamed Appendix in HPE report and 75(18.8%) patients had Normal Appendix in their HPE report.

332(83.0%) patients had Acute Appendicitis as final diagnosis and 68(17.0%) patients were with Normal Appendix as the final diagnosis.

Our study showed that in Acute Appendicitis, 282(84.9%) patients were with Elevated TLC and 50(15.1%) patients were with Normal TLC. In the Normal Appendix, 45(66.2%) patients were with Elevated TLC and 23(33.8%) patients had Normal TLC. The result has been shown in Table 1.

In Acute Appendicitis, 233(84.4%) patients had Caecal gurgling. In the Normal Appendix, 73(58.9%) patients had Cecal gurgling. The result has been shown in Table 2.

In our study, 347(86.8%) patients had Intraoperative Inflamed Appendix and 53(13.3%) patients were with Normal Appendix as shown in Table 3.

In our study, we have found that 325(81.3%) patients had Inflamed Appendix in HPE report and 75(18.8%) patients have Normal Appendix in their HPE report and the final result has been shown in Table 4.

DISCUSSION

In our study, we found among 332 patients (who were diagnosed with acute appendicitis in HPE report)

Table 1: Association between TLC: Final diagnosis

Final diagnosis			
TLC	Acute appendicitis	Normal appendix	Total
Elevated	282	45	327
Normal	50	23	73
Total	332	68	400

Chi-square: 13.3185; P: 0.0002. Odds Ratio: 2.8827 (1.6051, 5.1771)

Table 2: Association between cecal gurgling: Final diagnosis

Final diagnosis			
Cecal gurgling	Acute appendicitis	Normal appendix	Total
Absent	43	51	94
Present	233	73	306
Total	276	124	400

Chi-square: 31.0671; P: <0.0001, Odds Ratio: 0.2642 (0.1629, 0.4284)

Table 3: Association between intra-op finding: Final diagnosis

Final diagnosis			
Intra-op finding	Acute appendicitis	Normal appendix	Total
Inflamed appendix	318	29	347
Normal appendix	14	39	53
Total	332	68	400

Chi-square: 138.6375; P: <0.0001, Odds Ratio: 30.5468 (14.8785, 62.7150)

Table 4: Association between HPE FINDINGS: Final diagnosis

Final diagnosis			
HPE findings	Acute appendicitis	Normal appendix	Total
Inflamed Appendix	325	0	325
Normal Appendix	7	68	75
Total	332	68	400

Chi-square: 355.0201; P: <0.0001

282 patients were with elevated TLC and 50 patients were with normal TLC. The result was statistically significant(P=0.0002) as in Table 1.

Kamran et al.,⁸ showed in their study that the sensitivity and specificity of TLC as calculated in their study was 76.5% and 73.7%, respectively, while positive predictive value is 92.5%. TLC although not a diagnostic criteria for acute appendicitis but still was helpful investigation in decision making.

Ainippully et al.,⁹ had shown in their study that TLC proved more useful if it was performed within 24 h of the onset of symptoms and was relevant only if antibiotics have not been administered.

In our study, we found that in Acute Appendicitis, 233(84.4%) patients had Cecal gurgling. In the Normal Appendix, 73(58.9%) patients had Cecal gurgling. Association of Cecal gurgling versus Final Diagnosis was statistically significant (P<0.0001) as in Table 2.

MH Abbas et al.,¹¹ found in their studies that altered Alvarado scoring system in which an additional point of caecal gurgling was introduced, was slightly better in the diagnosis of acute appendicitis especially in equivocal patients (score 5-6). The rate of negative appendectomy in their study was 10.8%.

In this study, in Acute Appendicitis, 318(95.8%) patients had Inflamed Appendix and 14(4.2%) patients had Normal Appendix as per intra-operative findings. In the Normal Appendix, 29(42.6%) patients had Inflamed Appendix and 39(57.4%) patients were Normal Appendix as per intra-operative findings. Association of Intra-Op Finding versus The final diagnosis was statistically significant (P<0.0001) as in Table 3.

In Acute Appendicitis, 325(95.5%) patients had Inflamed Appendix and 7(2.1%) patients had Normal Appendix in HPE findings. In the Normal Appendix, 68(100.0%) patients had Normal Appendix. Association of HPE FINDINGS versus The final diagnosis was statistically significant (P<0.0001) as in Table 4.

In Acute Appendicitis, the mean MASS (mean±SD) of patients was 5.6476±.5914. In the Normal Appendix, the mean MASS (mean±SD) of patients was 4.5441±.8364. Difference of mean MASS with both Final Diagnosis was statistically significant (P<0.0001). In Inflamed Appendix, 261(80.3%) patients had Cecal gurgling. In Normal Appendix, 45(60.0%) patients had Caecal gurgling. Association of Cecal gurgling versus HPE FINDINGS was statistically significant (P<0.0001). In Acute Appendicitis, 312(94.0%) patient were ≥5 and 20(6.0%) patient were <5. In Normal Appendix, 22(32.4%) patient were ≥5 and

46(67.6%) patient were <5. Association of MASS versus Final Diagnosis was statistically significant ($P < 0.0001$).

Al Qahtani et al.,¹² found that the diagnosis of acute appendicitis remains mainly clinical, Alvarado score can be recommended as a helpful tool for the admission criteria and further management in order to reduce unnecessary admissions and to reduce the morbidity and mortality of acute appendicitis. The rate of negative appendectomy in their study was 12.5%.

In this study, we found that the modified Alvarado score was more in Acute Appendix compared to the Normal Appendix which was statistically significant. Cecal gurgling incorporating into Modified Alvarado score the Sensitivity was 94.0; Specificity was 67.6; Positive Predictive Value was 93.4; Negative Predictive Value was 69.7 and Accuracy was 95.0.

Sanjay et al.,¹³ found in their study that for MAS the sensitivity was 98.44%, specificity 94.4%.

Limitations of the study

In spite of every sincere effort our study has lacunae.

The notable short comings of this study are:

1. The sample size was small. Only 400 cases are not sufficient for this kind of study.
2. The study has been done in a single center.
3. The study was carried out in a tertiary care hospital, so hospital bias cannot be ruled out.

CONCLUSION

Keeping in view the results of the present study, we conclude that caecal gurgling is an important positive clinical finding in the diagnosis of acute appendicitis. Incorporating caecal gurgling into Modified Alvarado Scoring System, especially in the patients with a MAS of 5–6 where there is a dilemma whether to operate the patient or not, it aids in our decision making and thus helps to reduce the rate of negative appendectomies.

ACKNOWLEDGMENTS

I would like to express my gratitude to my guide, co guide, all respected seniors and beloved juniors of our department, my batchmates and faculties of pathology

and community medicine department. I am also thankful to the ethical committee, the patients and the patient party, the nursing staffs. Last but not the least i would like to mention my parents and my wife, without their constant support and inspiration this work could not be completed.

REFERENCES

1. Stephens PL and Mazzucco JJ. Comparison of ultrasound and Alvarado score for diagnosis of acute appendicitis. *Conn Med.* 1999;63(3):137-140.
2. Pal KM and Khan A. Appendicitis: A continuing challenge. *J Pak Med Assoc.* 1998;48(7):189-192.
3. Sonderegger MA. Diagnosis of Inflammatory Disease in Acute Abdomen; MD Thesis Paper, No. 777; 1939.
4. Drinkovic N. Age distribution and clinical characteristics in acute appendicitis. *Vojnosanit Pregl.* 1991;48(2):115-119.
5. Khan AH, Azhar MZ, Role of Alvarado score to minimise the rate of negative appendectomy without risk of perforation. *J Surg Pak.* 2007;12:93-97.
6. Barlas S, Günerhan Y, Palanci Y, İşler B and Çağlayan K. Epidemiological and demographic features of appendicitis and influences of several environmental factors. *Turk J Trauma Emerg Surg.* 2010;16(1):38-42.
7. Damanik B, Fikri E and Nasution IP. Relation between fiber diet and appendicitis incidence in children at H. Adam Malik central hospital. *Bali Med J.* 2016;5(2):268-271.
8. Kamran H, Naveed D, Nazir A, Hameed M, Ahmed M and Khan U. Role of total leukocyte count in diagnosis of acute appendicitis. *J Ayub Med Coll Abbottabad.* 2008;20(3):70-71.
9. Ainippully AM, Narayanan SK, Vysakh CN, Preeth VA and Somnath P. Role of total leukocyte count in diagnosis of pediatric acute appendicitis: Effect of antibiotic administration and duration of onset of symptoms. *Int Surg J.* 2018;5(1):296-299. <http://dx.doi.org/10.18203/2349-2902.isj20175914>
10. Yildirim O, Solak C, Koçer B, Unal B, Karabeyoğlu M, Bozkurt B, et al. The Role of serum inflammatory markers in acute appendicitis and their success in preventing negative laparotomy. *J Invest Surg.* 2016;19(6):345-352. <http://dx.doi.org/10.1080/08941930600985686>
11. Abbas MH, Altered alvarado score; role of caecal gurgling as a new sign in diagnosis of acute appendicitis. *J Sheikh Zayed Med Coll.* 2012;3(4):359-361.
12. Al Qahtani HH and Muhammad AA. Alvarado score as an admission criterion for suspected appendicitis in adults. *Saudi J Gastroenterol.* 2004;10(2):86-91.
13. Kumar SP, Karthik P, Karan RR. Determination of sensitivity and specificity of modified Alvarado score and ultrasonography in patients with acute appendicitis. *Int J Sci Res.* 2019;8(12):76-78.

Authors Contribution:

SD- Manuscript preparation, Data collection, Literature search; **SSK**- Conceptualized the study, literature search, data analysis and interpretation, revision of manuscript; **SR**- Literature search, prepared first draft; **KK**- Concept and design of the Study, Reviewed the literature and revision of manuscript, literature search

Work attributed to:

This work is attributed to our respected teachers and our beloved family members.

Orcid ID:

Dr. Susavan Das - <https://orcid.org/0000-0002-0001-677X>
 Dr. Sayak Roy - <https://orcid.org/0000-0003-0498-8194>
 Dr. Shib Shankar Kuri - <https://orcid.org/0000-0003-2212-6571>
 Dr. Kanchan Kundu - <https://orcid.org/0000-0002-6300-600X>

Source of Support: None, **Conflicts of Interest:** None.