

Prevalence and nature of psychiatric disorders in geriatric people attending psychiatry outpatient department

Poudel R¹, Belbase M²

1. Assistant Professor, Department of Psychiatry, Nepalgunj Medical College, Kohalpur, Nepal 2. Associate Professor, Department of Psychiatry, Nepalgunj Medical College, Kohalpur, Nepal

E-mail *Corresponding author : reet.poudel@gmail.com

Abstract

Introduction: Geriatric population is growing faster than all younger age groups. More than 20% of the geriatric people suffer from a mental or neurological disorder. Various studies have been conducted in Nepal related to mental and neurological disorders in geriatric people including epidemiological survey, outpatient department (OPD) based studies and inpatient studies. The aim of the present study was to determine the prevalence and nature of psychiatric disorders in geriatric subjects

Material And Method: Geriatric subjects attending outpatient department of psychiatry, Nepalgunj Medical College Teaching Hospital (NGMCTH), Kohalpur, Nepal from January to August 2018 (8 months) were enrolled for the study. Ethical clearance was obtained from the Institutional Review Committee of the NGMCTH and written consent was taken from the subjects or next to kin. Socio-demographic profile and other details of the subjects were obtained. Diagnosis was made using the ICD 10. Data was analyzed using Statistical Package for the Social Sciences (SPSS) version 21.0 for Windows.

Results: The prevalence of psychiatric disorders in geriatric subjects was 6.25%. The mean age of the subjects was 68.6 ± 7.39 years. Female constituted 51.1% of the subjects while male 48.9%. Category wise, mood disorders was the most common diagnosis (44.9%) followed by organic disorders (20.2%) and neurotic disorders (12.9%) respectively. Mood disorders and organic disorders were significantly associated with the age of the subjects ($p < 0.05$). Disorder wise, depression was the most common diagnosis (39.9%) followed by dementia (12.4%) and anxiety disorder (10.1%) respectively. Dementia and delirium were the only disorders significantly associated with the age of the subject ($p = 0.01$). There was no statistically significant association of individual disorders with sex of the subjects.

Conclusion: Depression, dementia and anxiety disorder was the most common diagnosis in geriatric people.

Keywords: Geriatric, Elderly, Psychiatric Disorder, Nepal

INTRODUCTION

Worldwide, geriatric population (60 years and above) is growing faster than all younger age groups. Compared to 2017, the number of persons aged 60 or above is expected to be more than double by 2050 and more than triple by 2100, rising from 962 million in 2017 to 2.1 billion in 2050 and 3.1 billion in 2100.¹ Similar to the global scenario, geriatric population has been increasing in Nepal since 1991, and in 2011

it constituted 8.13% of the total population of Nepal. It is expected to increase by 27.08% in 2021 and again by 25.23% in 2031 to comprise 10.20% of the total national population.²

The prevalence of mental disorders is high among the elderly.³⁻⁵ More than 20% of the geriatric people suffer from a mental or neurological disorder (excluding headache disorders) and 6.6% of all disability among them is attributed to mental and neurological

disorders. Dementia and depression are the most common mental and neurological disorders in this age group, which affect approximately 5% and 7% of the world's older population, respectively.⁶ As the geriatric population is rising all over the world, it expected to increase the burden of mental and neurological disorders in this age group in the coming future.

Various studies have been conducted in Nepal related to mental and neurological disorders in geriatric people including epidemiological survey⁷, outpatient department (OPD) based studies⁸⁻¹¹ and inpatient studies.¹² The aim of the present study was to determine the prevalence and nature of psychiatric disorders in geriatric subjects attending psychiatry OPD and to examine the relationship of various psychiatric disorders with the age and sex of the patient.

MATERIAL AND METHOD

The present study was conducted at Department of Psychiatry, Nepalgunj Medical College Teaching Hospital (NGMCTH), Kohalpur, Nepal. The duration of the study was eight months, from January to August 2018. Ethical clearance was obtained from the Institutional Review Committee of the NGMCTH. Subjects aged 60 years and above visiting the psychiatry OPD were explained about the study and written consent was taken from them or next to kin. Socio-demographic profile and other details of the subjects were obtained. Psychiatry diagnosis was made using the ICD 10.¹³ Data was analyzed using Statistical Package for the Social Sciences (SPSS) version 21.0 for Windows.

RESULT

During the period of 8 months, 2847 subjects visited psychiatry OPD among which 178 were in geriatric age group. The prevalence of geriatric subjects attending psychiatry OPD was 6.25%. The mean age of the subjects was 68.6 ± 7.39 years. Table 1 shows the socio-demographic characteristics of the subjects. Female constituted 51.1% (n=91) of the subjects while male 48.9% (n=87). Majority of the subjects were married (75.3%), Hindu (95.6%), from rural domicile (53.9%) and with no education (75.3%).

Table 1: Subjects Respondents' Socio-demographic Characteristics (n=178)

Characteristics		Frequency	Percentage
Sex	Male	87	48.9%
	Female	91	51.1%
Marital status	Never married	4	2.2%
	Married	134	75.3%
	Widow/Widower	37	20.8%
	Separated/Divorcee	3	1.7%
Place of Residence	Urban	80	44.9%
	Rural	96	53.9%
	India	2	1.1%
Religion	Hindu	170	95.6%
	Muslim	4	2.2%
	Christian	4	2.2%
Education	No education	134	75.3%
	Some education	44	24.7%

Table 2 shows the distribution of subjects based on age and sex. Category wise, mood disorders was the most common diagnosis (44.9%) followed by organic disorders (20.2%) and neurotic disorders (12.9%) respectively. Mood disorders and organic disorders were significantly associated with the age of the subjects (p<0.05). Disorder wise, depression was the most common diagnosis (39.9%) followed by dementia (12.4%) and anxiety disorder (10.1%) respectively. Dementia and delirium were the only disorders significantly associated with the age of the subject (p=0.01).

Table 3 shows distribution of subjects according to sex. There was no statistically significant association of individual disorders with sex of the subject.

Table 2: Distribution of Subjects According to Diagnosis & Age

Category	Diagnosis	Age range in years(N)					Total (N,%)	p value
		60-64	65-69	70-74	75-79	≥80		
Organic disorders	Dementia	4	5	2	4	7	22(12.4)	0.001*
	Delirium	0	0	3	3	2	8(4.5)	0.001*
	Organic Psychosis	1	2	3	0	0	6(3.4)	0.981
	Total	5	7	8	7	9	36(20.2)	0.000*
Substance Use Disorders	Alcohol use disorder	6	2	1	2	0	11(6.2)	0.230
Psychotic Disorders	Schizophrenia	2	3	3	0	0	8(4.5)	0.442
	Psychosis NOS	2	4	3	0	0	9(5.1)	0.569
	Total	4	7	6	0	0	17(9.6)	0.333
Mood Disorders	Depression	31	19	12	5	4	71(39.9)	0.05
	Mania	1	0	1	1	0	3(1.7)	0.626
	BPAD	0	3	2	0	1	6(3.4)	0.419
	Total	32	22	15	6	5	80(44.9)	0.020*
Neurotic Disorders	Anxiety Disorder	4	2	8	3	1	18(10.1)	0.250
	Somatoform Disorder	1	2	1	0	0	4(2.2)	0.479
	OCD	1	0	0	0	0	1(0.6)	0.244
	Total	6	4	9	3	1	23(12.9)	0.645
Other Disorders	Epilepsy	2	1	1	1	1	6(3.4)	0.452
	ISH	2	0	1	0	0	3(1.7)	0.214
	Primary Headache	1	1	0	0	0	2(1.1)	0.328
	Total	5	2	2	1	1	11(6.2)	0.598
Grand Total		58(32.6)	44(24.7)	41(23)	19(10.7)	16(9)	178(100)	

Table 3: Distribution Of Subjects According to Diagnosis and Sex

Category	Diagnosis	Sex		Total	p value
		Male	Female		
Organic disorders	Dementia	9	13	22	0.427
	Delirium	5	3	8	0.433
	Organic Psychosis	3	3	6	0.956
	Total	17	19	36	0.825
Substance Use Disorders	Alcohol use disorder	9	2	11	0.103
Psychotic Disorders	Schizophrenia	4	4	8	0.948
	Psychosis NOS	4	5	9	0.786
	Total	8	9	17	0.876
Mood Disorders	Depression	34	37	71	0.831
	Mania	2	1	3	0.537
	BPAD	5	1	6	0.087
	Total	41	39	80	0.570
Neurotic Disorders	Anxiety Disorder	7	11	18	0.374
	Somatoform Disorder	2	2	4	0.964
	OCD	1	0	1	0.308
	Total	10	13	23	0.581
Other Disorders	Epilepsy	2	4	6	0.441
	ISH	0	3	3	0.089
	Primary Headache	0	2	2	0.166
	Total	2	9	11	0.036*
Grand Total		87(48.88)	91(51.12)	178	

DISCUSSION:

Prevalence, age-sex distribution and socio-demographic profile:

Prevalence of geriatric subjects attending psychiatry OPD was 6.25%. Aich et al in an outpatient study reported an almost similar prevalence of 6.9%¹⁰. Nepalese OPD based studies related to psychiatry morbidity have reported 60 years and above subjects consisting of 7.6% to 11% of the total OPD attendance.¹⁴⁻¹⁶ In our study, male patients contributed 51.12% and female 48.18%. Male preponderance was observed by Aich et al¹⁰ (54.5%) whereas female preponderance was reported by Shakya (54%)¹¹ and Nepal et al (51%)¹⁷ in geriatric subjects visiting OPD. None of the psychiatric disorders were significantly associated with the sex of the subject. Majority of the patients (58%) were below 70 years of age. Similar findings have been reported in other OPD based studies from Nepal.^{10,11,17}

Psychiatric morbidity:

In our study mood disorder was the most common diagnosis category (44.9%). In OPD based studies from Nepal, mood disorders has been reported in the range of 19.5%¹⁰ to 36.7%.^{10,17} Shakya¹¹ reported a much higher prevalence of mood disorders of 46% but the sample consisted of subjects aged above 55 years while we included subjects 60 years and above. Study from India by Singh et al¹⁸ reported mood disorders as the most common diagnosis (48.07%) in geriatric subjects which is in consistent with our finding. Disorder wise depression was the most common diagnosis (39.9%). Prevalence of depression among elderly was found to be 49.4% in a community based study.¹⁹ Aich et al¹⁰ reported depression in 15.2% of geriatric subjects while life time prevalence of major depression in geriatric psychiatric patients was found to be 26.5% by Ritchie et al in French population.⁵ Our finding of depression as the most common diagnosis is in agreement with many earlier studies from Nepal.^{11,17}

Dementia (12.4%) was the second most common diagnosis. Studies from Nepal have shown dementia in range of 6%¹¹ to 8.9%¹⁰ in geriatric OPD visiting patients. Organic mental disorders as a group was observed in 20.2%, the value of which was similar to that reported by Koirala et al (28%).²⁰ Our study revealed anxiety disorder

as the third most common diagnosis with 10.1% geriatric patients diagnosed with it. Koirala et al and Aich et al reported a similar low prevalence of anxiety disorder of 8% and 9.7% respectively. Shakya reported a higher prevalence of 22% but the disorders included not only anxiety disorder exclusively, but also phobia and obsessive compulsive disorder. Neurotic disorders as a group was observed in 12.9% of the subjects which is similar to that observed by Nepal et al (13.8%)¹⁷. Singh et al¹⁸ reported 15.47% of geriatric subjects diagnosed with neurotic disorders which is consistent with our study. Aditya et al²¹ in a study in Kathmandu found higher prevalence of anxiety disorders (76.1%) in geriatric patients. A different methodology employed for the study and subjects aged 65 years and more suffering from some medical illness enrolled for the study might have resulted in higher prevalence of anxiety disorder in the said study.

Psychotic disorders were observed in 9.6% of the subjects. Shakya (6%) and Nepal et al. (6.2%) reported a similar low prevalence of psychotic disorders in elderly whereas Aich et al¹⁰ reported a higher prevalence (26.4%) which may have resulted due to difference in the geographical location of the conducted studies. Alcohol use disorder (AUD) was diagnosed in 6.2% of the subjects. Nepalese OPD based studies have reported AUD in range of 10.5% to 14%.^{10,11,17,22}

In our study prevalence of BPAD (including first episode mania) was 5.06%. Reported prevalence of mania amongst geriatric people is found to be low (1-2%) in both in-patient as well as community prevalence studies from western countries.^{5,23} Aich et al¹⁰ reported prevalence of mania as 4.3% while Shakya¹¹ reported prevalence of BPAD as 7% which is in agreement with our finding. We also reported prevalence of other disorders which frequently present to psychiatry OPD including epilepsy (3.4%), intentional self harm or ISH (1.7%) and primary headache (1.1%). Nepal et al reported prevalence of headache as 1.90% which is similar to our study. Aich et al¹⁰ reported a much higher prevalence of chronic headache (12.8%). in this age group. This discrepancy of prevalence of primary headache may be due to geographical location of the institution and due to increased awareness of other departments in

referring headache to psychiatry. ISH has been reported in the range of 0.95%¹⁷ to 2%¹¹ in earlier Nepalese studies which is similar to our study. Depression, dementia and anxiety disorder was the most common diagnosis. Category wise, mood disorders and organic disorders were significantly associated with the age of the subjects ($p < 0.05$). Dementia and delirium were the only individual disorders significantly associated with the age of the subject ($p = 0.01$). In our study, we tried to determine the prevalence of psychiatric disorders in geriatric people attending psychiatry OPD and analyzed the relationship of psychiatric disorders with the age and sex of the subject. As this study was conducted in OPD attending subjects, there were some limitations that need to be addressed. First, OPD based finding cannot not be generalized to the population of the community so more community based research on this topic is warranted. Second, geriatric subjects were not accessed for any medical or surgical comorbidity which might have increased the presentation of psychiatric disorders like anxiety and depression.

CONCLUSION:

The present study found the prevalence of geriatric subjects attending psychiatry OPD to be 6.25%. Depression, dementia and anxiety disorder was the most common diagnosis in geriatric people visiting psychiatry OPD.. Category wise, mood disorders and organic disorders were significantly associated with the age of the subjects ($p < 0.05$). Dementia and delirium were the only individual disorders significantly associated with the age of the subject ($p = 0.01$). There was no statistically significant association of individual disorders with sex of the subject.

ACKNOWLEDGEMENT: None

CONFLICT OF INTEREST: None

REFERENCES:

1. *World Population Prospects: The 2017 Revision, Key Findings and Advance Tables. Working Paper No. ESA/P/WP/248. United Nations, Department of Economic and Social Affairs, Population Division; 2017.*
2. *2015 Statistical Year Book Nepal. Government of Nepal National Planning Commission Secretariat Central Bureau of Statistics; 2016.*
3. Gum AM, King-Kallimanis B, Kohn R. *Prevalence of mood, anxiety, and substance-abuse disorders for older Americans in the national comorbidity survey-replication. The American Journal of Geriatric Psychiatry. 2009 Sep;17(9):769-81.*
4. Prévaille M, Boyer R, Grenier S, Dubé M, Voyer P, Puntis R, et al. *The epidemiology of psychiatric disorders in Quebec's older adult population. Canadian Journal of Psychiatry. 2008 Dec;53(12):822-32.*
5. Ritchie K, Artero S, Beluche I, Ancelin M, Mann A, Dupuy A, et al. *Prevalence of DSM-IV psychiatric disorder in the French elderly population. The British Journal of Psychiatry. 2004 Feb;184:147-52.*
6. World Health Organization. *Mental health of older adults [Internet]. [cited 2018 Jul 11]. Available from: <https://www.who.int/news-room/fact-sheets/detail/mental-health-of-older-adults>*
7. Subedi S, Tausig M, Subedi J, Broughton CL, Williams-Blangero S. *Mental illness and disability among elders in developing countries: the case of Nepal. Journal of Aging and Health. 2004 Feb;16(1):71-87.*
8. Khattri JB, Nepal MK. *Study of depression among geriatric population in Nepal. Nepal Medical College Journal. 2006 Dec;8(4):220-3.*
9. Ghimire S, Baral BK, Pokhrel BR, Pokhrel A, Acharya A, Amatya D, et al. *Depression, malnutrition, and health-related quality of life among Nepali older patients. BMC Geriatrics. 2018 Aug 24;18(1):191.*
10. Aich TK, Shah S, Subedi S. *Pattern of Neuropsychiatric Illnesses in Geriatric Population: An Outpatient Study Report. Journal of Psychiatrists' Association of Nepal. 4(1):12-9.*
11. Shakya DR. *Psychiatric morbidities of elderly psychiatry out-patients in a tertiary-care hospital. Journal of College of Medical Sciences-Nepal. 2011;7(4):1-8.*
12. Aich TK, Dhungana M, Muthuswamy R. *Pattern of neuropsychiatric illnesses in older age group population: an inpatient study report from Nepal. Indian Journal of Psychiatry. 2012 Jan;54(1):23-31.*
13. *The ICD-10 Classification of Mental and Behavioral Disorders Diagnostic Criteria for Research. Geneva: World Health Organization; 1993.*
14. Shrestha MR, Pradhan S, Sharma S. *Morbidity pattern of psychiatric disorders in patient seeking treatment in psychiatric OPD of private tertiary care hospital. Postgraduate medical journal of NAMS. 2011;11(1):28-33.*
15. Koirala NR, Yadav R, Das AK, Poudel J, Bhagat SK. *A study of socio-demographic and diagnostic profile of patients attending the psychiatric out-patient department of Nobel Medical College, Biratnagar. Journal of Nobel Medical College. 2012;1(2):45-9.*
16. Khattri JB, Godar ST, Thapa P, Ramesh K, Chakraborty PK, Thapa BB. *Socio-demographic*

- characteristics and diagnostic profile of patients attending psychiatric OPD of a private hospital in western region of Nepal. *Nepal Journal of Medical Sciences*. 2012;1(1):15-8.
17. Nepal S, Sapkota N, Kumar R, Deo BK, Mishra S. Psychiatric Disorders in Elderly Patients attending OPD of Tertiary Care Centre in Eastern region of Nepal. *Journal of Psychiatrists' Association of Nepal*. 2016;5(1):221-6.
 18. Singh GP, Chavan BS, Arun P, Lobraj, Sidana A. Geriatric out-patients with psychiatric illnesses in a teaching hospital setting - a retrospective study. *Indian Journal of Psychiatry*. 2004 Apr;46(2):140-3.
 19. Subedi S, Shrestha P, Thapa D. Study Of Depression in Elderly: Prevalence and Factors Associated. *Journal of Psychiatrists' Association of Nepal*. 2018;7(2):16-23.
 20. Koirala NR, Mahat P, Nepal MK, Ojha SP, Sinha UK, Pokhrel A, et al. Geriatric psychiatry: socio-demographic characteristics and diagnostic profile among senior citizens attending the psychiatric outpatient department of a tertiary health care facility in Nepal. *Journal of Institute of Medicine*. 2000;22(3).
 21. Kumar, A, Sharma SR, Timilsina S, Giri S, Yadav V. High Prevalence of Depression and Anxiety Symptoms among Hospitalized Geriatric Medical Inpatients: A study from a Tertiary Level Hospital in Nepal. *University of Toronto Medical Journal*. 2010;88(1):32-5.
 22. Thapa P, Chakraborty PK, Khattri JB, Ramesh K, Sharma B. Psychiatric morbidity in elderly patients attending OPD of tertiary care centre in western region of Nepal. *Industrial Psychiatry Journal*. 2014;23(2):101-4.
 23. Draper B. The elderly admitted to a general hospital psychiatry ward. *The Australian and New Zealand Journal of Psychiatry*. 1994 Jun;28(2):288-97.