

# Pathological Laughing in a Patient with a Pontine Tumor

Gopi Nepal<sup>1</sup>, Pritam Gurung<sup>1</sup>, Anik Jha<sup>1</sup>, Durga Khadka<sup>2</sup>, Resha Shrestha<sup>1</sup>, Basant Pant<sup>1</sup>


<sup>1</sup>Department of Neurosurgery, Annapurna Neurological Institute and Allied Sciences, Maitighar, Kathmandu, Nepal

<sup>2</sup>Department of Psychiatry, Annapurna Neurological Institute and Allied Sciences, Maitighar, Kathmandu, Nepal

Date of submission : 18<sup>th</sup> Feb 2023

Date of acceptance: 15<sup>th</sup> May 2023

Date of publication: 15<sup>th</sup> July 2023



## Abstract

An 18-year-old man presented with the complaints of occasional headache and limb weakness associated with slurring of speech and purposeless laughing. Magnetic resonance imaging showed a diffuse altered signal intensity area involving the pons with asymmetrical expansion. The patient underwent surgical resection. Histology was suggestive of a WHO grade IV glioblastoma. Pathological laughing in a patient with a pontine lesion is a very rare presentation. Evidence for a mechanism of action underpinning how pontine lesions can cause such behavioral changes through the disruption of a network of cerebro-ponto-cerebellar pathways is discussed.

**Key words:** Pathological laughing, Pons, tumor

## Introduction

Pathological laughing and crying (PLC) can be described as uncontrollable episodes of laughing or crying that are triggered by a stimulus that would not normally cause either response.<sup>1</sup> Patients with a lesion in the pons mainly present with features of motor deficits, and associated cranial nerve palsies. However, pathological laughter in a patient with a pontine lesion is a very rare presentation. The disruption of a network of cerebro-ponto-cerebellar pathways has been considered to be a possible mechanism for such behavior. We report a case of pathological laughing in a patient with pontine tumor.

## Case report

An 18 year-old man presented with a headache for 1 month and limb weakness for 10 days. He also had slurring of speech and purposeless laughing while talking over the same duration. On examination, motor power in the lower limb was 4/5, with increased tone and exaggerated deep tendon reflexes. Ankle clonus was present. Hand grip was weak in both hands (Right > Left). A sensory exam appeared to be intact.

Magnetic resonance imaging (MRI) showed a diffuse altered signal intensity area involving the pons with its asymmetrical expansion predominantly on the left aspect measuring 29 x 32 x 39 mm (Figure 1A-1B). The lesion was causing dorsal displacement of the floor of the fourth ventricle, with compression of the middle cerebellar peduncle. This was towards the left side, was abutting both the basilar artery and its anterior displacement with normal flow voids. The post contrast image showed thick, irregular, and heterogeneous enhancement within the expanded pons (Figure 2 A-2B). Magnetic resonance spectroscopy showed a decrease in NAA and an elevated choline/ creatine ratio. The patient was diagnosed with pontine tumor and underwent a left retrosigmoid craniectomy, and gross total excision of the tumor was achieved. Postoperative imaging showed complete resection of the tumor. Histopathological examination was suggestive of glioblastoma (Figure 3). A customized immunohistochemistry panel was advised, which revealed immunoreactive Olig-2, Ki-67, ATRX, P53 and H3K27M.

The patient's postoperative recovery was uneventful with mild hemiparesis and facial palsy. The patient was discharged on the tenth postoperative day with satisfactory outcomes and a Karnofsky performance score (KPS) of

### Access this article online

Website: <https://www.nepjol.info/index.php/NJN>

DOI: <https://doi.org/10.3126/njn.v20i2.52611>

### HOW TO CITE

Nepal G, Gurung P, Jha A, Khadka D, Shrestha R, Pant B. Pathological Laughing in a Patient with a Pontine Tumor. NJNS. 2023;20(2):75-78.



### Address for correspondence:

Gopi Nepal

FCPS Resident

Annapurna Neurological Institute And Allied Sciences

Maitighar, Kathmandu

Tel number: 977-9860525799

Email: [gopinepal1000@gmail.com](mailto:gopinepal1000@gmail.com)

Copyright © 2023 Nepalese Society of Neurosurgeons (NESON)

ISSN: 1813-1948 (Print), 1813-1956 (Online)




This work is licensed under a Creative Commons Attribution-Non Commercial 4.0 International License.

50%. Postoperatively, the patient was then scheduled for referral to a cancer center for further treatment. However,


he unfortunately passed away in a local hospital while receiving treatment for pneumonia.



*Figure 1: Magnetic resonance imaging (T1 axial and sagittal) shows diffuse altered signal intensity area involving the pons with its asymmetrical expansion predominantly on the left aspect measuring 29 x 32 x 39 mm. The lesion is causing dorsal displacement of floor of the fourth ventricle with compression of the middle cerebellar peduncle more on the left side, and is abutting the basilar artery and its anterior displacement with normal flow voids.*



*Figure 2: Magnetic resonance imaging contrast (T1 coronal) shows thick irregular and heterogeneous enhancement within the expanded pons.*



*Figure 3: Histopathological examination showing increased cellularity and the cells are arranged in diffuse pattern in a fibrillary background. The cells are moderate to markedly pleomorphic and hyperchromatic with abundant eosinophilic cytoplasm. The nuclei are round, oval and elongated. Microvascular proliferation, few gemistocytes, necrosis in a tiny focal area and hemorrhage are also seen. (Magnification 10 X)*



Figure 4: Post operative CT scan of head revealed gross total removal of tumor.

### Discussion

Pathological laughing has been reported across literature in cases of tumor<sup>1,2</sup>, infarction and abscess<sup>3</sup>. The exact mechanism is still poorly understood, with various pathological proposals. Oppenheim, Siemerling, and Wilson suggested that modulation is facilitated by direct corticobulbar pathways and in a linear top-down model.<sup>4,5</sup> They posited that PLC occurs when the voluntary control of the emotional expression fails due to bilateral lesions of the descending corticobulbar tracts. Parvizi et al, suggested that an intact relationship between the cerebral cortex and cerebellum is important for a normal regulation of emotional expression.<sup>6</sup> They propose therefore that problems with an exaggerated or contextually inappropriate emotional response result when the cerebellar modulation of these behaviors is impaired by a lowered emotional threshold, or by an incorrect, contextually inappropriate response. Wilson et al, suggested that a brainstem faciorespiratory center could act as a mechanism of this phenomenon.<sup>5</sup> This center was presumed to be under the control of higher centers, and pathological laughter was thought to ensue when they were incapable of exerting their inhibitory influence on the faciorespiratory center.

Structures such as the pre frontal cortex, anterior cingulate cortex, internal capsule, thalamus, subthalamic nucleus, basis pontis, and cerebellar white and gray matter have been implicated in PLC, as revealed by many studies. The basis pontis stands out as the only identified site where a discrete lesion can cause PLC.<sup>7</sup>

Functional imaging studies have proved helpful in discovering the neural correlates for both pathological and normal regulation of emotional expression.<sup>8</sup> They used three types of fMRI experiment tasks, including facial recognition, semantic decision, and motor function to establish the response to non-specific stimuli. The patient showed consistently abnormal pontine activation while performing all tasks before treatment, which was not present for any of the controls. However, the exaggerated pontine activation was normalized after patients were treated with paroxetine, terminating the laughing episodes. Thus, their study suggested that serotonergic replacement decreases the aberrant activity in a circuit that involves the pons.

Various reports suggested that tricyclic antidepressants such as amitriptyline and nortriptyline and selective serotonin reuptake inhibitors (SSRIs) are effective in treating PLC.<sup>9,10</sup> Abnormal fMRI results disappearing with clinical improvement after treatment with paroxetine also suggested some SSRI efficacy in PLC.<sup>8</sup>

Pathological laughing was the only feature shown by the case reported. The patient's postoperative recovery was uneventful with mild hemiparesis and facial palsy. He had no purposeless laughing after the surgery. The patient was discharged with satisfactory outcome with a KPS of 50%. Postoperatively, the patient was scheduled for referral to a cancer center for further treatment. However, he suffered from pneumonia and passed away in another local hospital during the pneumonia treatment process.

### Conclusion

We have presented a very rare case of pathological laughing in a patient with a pontine tumor. Patients with the features of PLC are often underestimated. Therefore, these symptoms may serve as an early indicator for disease progression. An understanding of the mechanisms underlying these behavioral changes may help in providing both clinical and psycho-social care.

**Conflicts of interest:** None of the authors have potential conflicts of interest to be disclosed.

### References

1. Cantu RC, Drew JH. Pathological laughing and crying associated with a tumor ventral to the pons. Case report. *J Neurosurg.* 1966. doi:10.3171/jns.1966.24.6.1024
2. Hargrave DR, Mabbott DJ, Bouffet E. Pathological laughter and behavioural change in childhood pontine glioma. *J Neurooncol.* 2006. doi:10.1007/s11060-005-9034-8

3. Elyas AE, Bulters DO, Sparrow OC. Pathological laughter and crying in patients with pontine lesions: Case report. *J Neurosurg Pediatr.* 2011. doi:10.3171/2011.8.PEDS11265
4. Text-Book of Nervous Diseases for Physicians and Students. By Professor H. Oppenheim, of Berlin. Authorised translation by Alexander Bruce, M.D., F.R.C.P.E., LL.D. Fifth enlarged and improved edition. Edinburgh, 1911. 432 illustrations and 8 plates. *Pric. J Ment Sci.* 1911. doi:10.1192/bjp.57.237.376
5. Wilson SAK. Some Problems in Neurology: Pathological laughing and crying. *Psychopathol J Neurol.* 1924.
6. Parvizi J, Anderson SW, Martin CO, Damasio H, Damasio AR. Pathological laughter and crying: A link to the cerebellum. *Brain.* 2001. doi:10.1093/brain/124.9.1708
7. Parvizi J, Coburn KL, Shillcutt SD, Coffey CE, Lauterbach EC, Mendez MF. Neuroanatomy of pathological laughing and crying: A report of the american neuropsychiatric association committee on research. *J Neuropsychiatry Clin Neurosci.* 2009. doi:10.1176/jnp.2009.21.1.75
8. Kosaka H, Omata N, Omori M, et al. Abnormal pontine activation in pathological laughing as shown by functional magnetic resonance imaging. *J Neurol Neurosurg Psychiatry.* 2006. doi:10.1136/jnnp.2005.073288
9. Schiffer RB, Herndon RM, Rudick RA. Treatment of Pathologic Laughing and Weeping with Amitriptyline. *N Engl J Med.* 1985. doi:10.1056/nejm198506063122303
10. Burns A, Russell E, Stratton-Powell H, Tyrell P, O'Neill P, Baldwin R. Sertraline in stroke-associated lability of mood. *Int J Geriatr Psychiatry.* 1999. doi:10.1002/(SICI)1099-1166(199908)14:8<681::AID-GPS49>3.0.CO;2-Z