

BURDEN OF ENTEROPATHOGENS ASSOCIATED DIARRHEAL DISEASES IN CHILDREN HOSPITAL, NEPAL

Jeevan B. Sherchand*, Michiyo Yokoo**, Ojaswee Sherchand*, Arjun R. Pant***, Osamu Nakagomi**

*Tribhuvan University, Institute of Medicine, Public Health Research Laboratory and Microbiology, Kathmandu, Nepal.

**Division of Molecular Epidemiology Graduate School of Biomedical Sciences, Nagasaki University, Sakamoto, Nagasaki, Japan.

***Ministry of Health, Kanti children Hospital, Maharajgunj, Kathmandu, Nepal.

Abstract: Diarrheal disease caused by bacteria, parasites or viruses continues to be an important cause of morbidity and mortality among young children in developing countries. Methods currently used in public health laboratories do not allow for the identification of rotavirus, *Cyclospora* and pathogenic *E.coli* infection though they represent as an etiology in large proportion of patients with diarrhea, the possibility exists that a portion of the undiagnosed illness may be attributable to one or more of the above enteropathogens. In a view to determine the causative agents of diarrhoea, the current study described the various enteropathogens associated with diarrhoea in hospitalized children.

Stool samples were collected from children under 11 years of age who developed diarrhoea and were admitted to Kanti Children's Hospital between May to October 2007 and investigated in Tribhuvan University, Institute of Medicine, Health Research Laboratory; by using both the combination of microbiological and immunological tools (EIA for rotavirus detection, standard parasitological procedure for *Cyclospora* and other intestinal parasites, and selective culture method and serotyping were used to differentiate the species of bacteria).

A total of 440 diarrhoeal stool samples were collected and 285 (64.8%) enteropathogens were identified. The highest infection was due to intestinal parasites 104/285 (36.5%) followed by rotavirus 92/285 (32.3%); pathogenic bacteria 57/285 (20%) and *Cyclospora* 32/285 (11.2%). Among the pathogenic bacteria (20%) isolated, the predominant bacteria were *Shigella species* (36.8%); *Vibrio species* (26.3%); *Escherichia coli* (22.8%) and *Salmonella species* (14.03%) respectively.

Various enteropathogens responsible for diarrhoea especially rotavirus, different pathogenic bacteria and *Cyclospora* infection, which are not examined routinely in public health laboratories, were found in significant proportion as a cause of diarrhoeal illness in children. The infection was peak in children under 2 years of age and was highest in rainy season.

Key words: Burden; Diarrhea; Enteropathogens; Children; Nepal.

INTRODUCTION

Diarrheal disease caused by bacteria, parasites or viruses continues to be an important cause of morbidity and mortality among young children in developing countries¹. Cumulative data from epidemiological studies show that approximately 20–50% of cases are attributable to known bacterial or parasitic pathogens, which suggests that viruses may be responsible for the remainder². In fact, viral gastroenteritis has been found to be the second most common viral clinical entity in developed countries, following closely behind viral upper respiratory tract illness³⁻⁴.

Immense micro floras are implicated in diarrhoeal illness⁵⁻⁷. Some of these, including *Salmonella*, *Shigella* and *Vibrio spp.* are well recognized enteric pathogens. Rotavirus, *Campylobacter spp.* and enterotoxigenic *E. coli* (ETEC) also have been identified as important causes. Recently, certain strains of *Aeromonas hydrophila* and *Plesiomonas shigelloides* have been associated causally by some investigators⁸⁻¹⁰. The coccidian parasite *Cyclospora cayetanensis* is a newly recognized enteric pathogen causing prolonged diarrhea in humans¹¹⁻¹². It has been implicated as an important cause of diarrhoeal illness in the context of Nepal¹³⁻¹⁶.

Author for Correspondence: Dr. Jeevan Bahadur.Sherchand, Tribhuvan University, Institute of Medicine, Health Research Laboratory and Clinical Microbiology, P.O. Box: 13913; Kathmandu, Nepal.

Rotaviruses are reported to be the most common cause of severe childhood diarrhea worldwide¹⁷. Illness associated with rotavirus has also been observed in older children and adults. Rotaviral gastroenteritis may result in mortality for populations at risk such as infants and the elderly¹⁸. Rotavirus was found in 25-40% of children with diarrhea in urban Kathmandu valleys of Nepal¹⁹⁻²⁰.

In Nepal however, the major causative agents; rotavirus, *Cyclospora* and different species of pathogenic bacteria are not routinely examined though they are important causes of diarrhea in children. Therefore, this study is carried out to reveal the etiology of diarrhoeal illness in high risk population group especially young children less than 11 years of age using such unique tools.

MATERIALS AND METHODS

The study was carried in Kanti Children Hospital and Tribhuvan University Teaching Hospital, Public Health Research Laboratory, Nepal between May to October 2007 and included 440 hospitalized patients with acute diarrhea. From these hospitalized 440 children, stool specimens were collected and immediately transported to the laboratory and all stool were cultured for bacterial investigation within 2 hours of collection on Salmonella-Shigella agar (HI-media, UK), Mac Conkey agar (Oxoid, England) and TCBS agar (HI-media.) plates and incubated overnight at 37°C. They were also cultured in Selenite F broth and peptone broth (HI-media) and incubated up to 4 hours. When there was growth in broth, an inoculum was sub cultured on SS agar and TCBS plates. After culture, the stool specimens were examined for parasites and rotavirus antigen detection. Rota clone ELISA kits (Meridian Bioscience Inc.) were used to detect rotavirus and sucrose floatation concentration methods were used for coccidian parasites *cyclospora* and *cryptosporidium*. They were then confirmed by modified Ziehl-Neelson's technique.

In addition, a structure questionnaire was filled out for each patient covering demographic information, area of residence, information about any symptoms associated with the disease (fever, vomiting, dehydration status), type and duration of diarrhea and type of treatment received—if any—in the hospital. Acuter diarrhea was defined as abnormal fecal discharge characterized by frequent—at least three times per day—liquid or semi-liquid loose stools, accompanied by symptoms such as nausea, vomiting and fever and involving dehydration and electrolyte loss.

RESULTS

A total of 440 children with diarrhea were enrolled in which 248 (56.4%) were males and 192 (43.6%) were females.

Clinical symptoms in diarrhoeal patients:

Different clinical symptoms and other features such as duration of diarrhea, fever, vomiting, nausea and degree of dehydration and abdominal pain are depicted in table 1.

Table 1: Clinical symptoms and other features of diarrhoeal children

Clinical symptoms	No (%)
Diarrhoeal duration	
3-7 days	201(45.7)
7-10 days	89(20.5)
10—14 days	80(18.2)
>15 days	70(15.9)
Fever	197(44.8)
Vomiting	410(93.2)
Nausea	311(70.7)
Dehydration	
No	129(29.3)
Mild to moderate	254(57.7)
Severe	57(13)
Abdominal pain	279(63.4)

Distribution of enteropathogens

The distributions of enteropathogens in different age group are shown in table 2. The highest rate of intestinal parasites (30.9%), followed by rotavirus infection (20.9%) and pathogenic bacterial infection; 12.9%. The highest rate of rotavirus infection was found in age group between 0-2 years and the lowest rate was found age between 6-8 years of age, but the highest rate of bacterial infection was found age between 6-8 and the lowest was in 0-2 years of age. Intestinal parasitic infection was found highest among the age between 9-11 and the lowest was between 0-2 years of age. (Table 2).

Table 2: Age wise distribution of different enteropathogens

Age group	Total No	Rotavirus +ve(%)	Bacteria +ve(%)	Parasites +ve(%)
0-2	195	56(28.7)	15(7.7)	44(22.5)
3-5	103	24(23.3)	19(18.5)	28(27.2)
6-8	62	5(8.1)	12(19.4)	26(41.9)
9-11	80	7(8.8)	11(13.8)	38(47.5)
Total	440	92(20.9)	57(12.9)	136(30.9)

Intestinal parasites among diarrhoeal children:

In the study 136 (30.9%) diarrhoeal children were infected with intestinal parasites in which the highest parasitic infection rate was *giardia lamblia* (20.6%) and the lowest rate *cryptosporidium parvum* (2.9%). The reemerging coccidian parasites *Cyclospora cayentanensis* associated with diarrhoeal disease as depicted in table 3 was found 8.8% among the hospitalized children.

Table 3: Distribution of Intestinal Parasites

Parasites	No of Positive (%)
<i>Ascaris lumbricoides</i>	15(11.02)
<i>Trichuris trichiura</i>	11(8.08)
Hookworm	9(6.6)
<i>Hymenolepis nana</i>	11(8.08)
<i>Giardia lamblia</i>	28(20.6)
<i>Entamoeba histolytica</i>	21(15.4)
<i>Entamoeba coli</i>	21(15.4)
<i>Balantidium coli</i>	4(2.9)
<i>Cyclospora cayentanensis</i>	12(8.8)
<i>Cryptosporidium parvum</i>	4(2.9)
Total no. of parasites	136

Distribution of Pathogenic Bacteria:

Pathogenic bacteria were isolated in 57 (12.9%) cases as shown in figure 1 and table 4.




Figure 1: Distribution of bacterial isolates

Table 4: Distribution of bacterial pathogens (N=57)

Type of organism	No of Organism
Vibrio Cholerae 01	15(26.3%)
Ogawa	12
Hikojima	2
Inaba	1
Escherichia coli	13(22.8%)
EPEC	8
ETEC	2
EHEC	2
EIEC	1
Salmonella spp	8(14.03%)
S. paratyphi A	3
S. paratyphi B	2
S. paratyphimurium	2
S. typhi	1
Shigella spp	21(36.8%)
S. dysenteriae	14
S. boydii	5
S. flexneri	2
S. sonnei	0

Distribution of Rotavirus infection

Of the total 440 sample examined, 92 (20.9%) cases were found rotavirus positive. The highest rate was found in age group 0-2 years as shown in figure 2.




Figure 2: Age wise distribution of rotavirus infection

DISCUSSION

Diarrhoeal diseases are the main social problems in Nepal as in other developing countries in tropical zones²¹⁻²⁴. The present study which covered the peak diarrhoeal seasons of six months (from May to October 2007), was conducted to determine the prevalence of viral, parasitic and bacterial enteropathogens associated with diarrhea in hospitalized children under 11 years of age. Using a combination of traditional and molecular diagnostic techniques, 285 (64.8%) enteropathogens were detected in total of 440 patients with diarrhea studied.

A total of 136 (30.9%) intestinal parasites infestation children 104 cases (23.6%) of other than *Cyclospora* infestation were found in which protozoan dominated over helminthes. Among the helminthes, the highest prevalence was observed for *Ascatis lumbricoides* followed by *Trichuris trichiura* and *Hymenolepsis nana* and Hookworm. Similarly, *Giardia lamblia* was followed by *Entamoeba histolytica* *Balantidium coli* and *Entamoeba coli*. were found respectively among the protozoan. Coccidian parasite, *Cryptosporidium parvum* was detected in 2.9% cases. Despite the government policy for antihelminthic program and health education program launched at community level, the intestinal parasite infestation still remains the major cause of diarrheal diseases in children. This can be taken as the inadequate implementation of the program or can be said as the carelessness of people themselves as they take diarrheal disease as a minor disease, often being termed as social disease in many developing countries by different researchers. Coincidentally, parasitic infestation peaks during the warmer and rainy season reflecting poor drainage and waste management and also as the favorable condition for the disease transmission.

C. cayetanensis, an emerging pathogen was detected in 12(8.8%) of the total cases, with highest infection rate in the age group 6-8 years. Initial studies indicated that it causes diarrhea mainly in immunocompromised patients and prolonged diarrhea in expatriate populations in certain developing countries including Nepal²⁵. However, a subsequent study in Nepal suggested that it also causes diarrhea among the indigenous children¹³.

The 13% of bacterial pathogens were also found as significant cause of diarrhea in this study. Among them, *Shigella spp* (21 cases, 36.8%) was most frequently detected followed by *Vibrio spp* (15 cases, 26.3%), *Escherichia coli* (13 cases, 22.8%) and *Salmonella spp* (8 cases, 14.03%). The age group 6-8 years was found highly infected (19.4%). The highest prevalence of bacterial infection found in this age group can be explained as the children at this age group are more actively involved in playing outdoors and careless about their feeding and drinking habits, lacking in proper knowledge of health education and proper personal hygiene, etc. And immunity to combat bacterial infection is also not well developed at this age. Moreover, the prevalence of bacterial pathogen reported herein could be affected by the fact that the reliable data on exclusion of self-treated individuals could not be made available.

Rotavirus was the most prevalent viral etiologic agent and was detected in 20.9% of the hospitalized children. Our findings were supported by Sherchand and Haruki²⁶, who reported that approximately 30.6% of infants and young children under 5 years of age with diarrhea from hospitals were infected with rotavirus. The prevalence of rotavirus in Nepal as estimated by present study is comparable to that reported for other developing and developed countries, which ranges from 30-50%⁴. The predominance of rotavirus is clearly the reflection of age group- 60.9% were under the age group 0-2 years. The finding is in accordance with similar results found by other researchers^{19, 21, 27-31}. It appeared that infants below 4 months of age were initially protected to some extent by maternal antibodies against severe diarrhea due to rotavirus, and they seem to have acquired adequate immunity between 12 and 16 months of age. The greater risks of infants and young children in the period between 6 to 12 months with declined levels of maternal antibodies to rotavirus infection have been documented³²⁻³³.

Seasonal variation of enteropathogen infection indicated that the infection rate was highest in peak rainy season (July and August). In particular, parasitic infection was found highest followed by rotavirus, bacterial pathogen. This reveals that incidence of enteropathogen appears to be due to the heavy fecal contamination of drinking water, rapid and unplanned urbanization, improper disposal of wastes, poor sanitary facilities and health education, etc. In many parts of cities (Kathmandu, Lalitpur and Bhaktapur) in Kathmandu valley, drinking water supplies has been reported to be contaminated with sewage due to running of water pipe and sewage line closer together, water flowing intermittently causing negative pressure thus suckling fluids and air from leaky pipe lines³⁴. In developing countries, contamination of water may occur at the source, but may also occur at the time of collection and storage at home³⁵.

Though rotavirus infection peaks during winter seasons, this study showed the prevalence rate of rotavirus infection as 20.9% during the warmer months like May to October which is in accordance with the study that identified rotavirus more often in warmer months³⁶. The seasonal nature of rotavirus infection was not universal, and in countries within 10° of the equator, infection occurred year around³⁷. Rotavirus is a disease of later warm and early cool season but seasonality may vary year to year³⁸.

Cyclospora infection was found highest in June which was in accordance with the findings of other researchers^{16, 39}. The environmental conditions during this season are favorable for sporulation of cysts and thus help to maintain the chain of infection.

Often assignment of an etiologic agent role in specific cases of diarrhea was particularly difficult because more than one potential pathogens were isolated from the same stool specimen in 8.2% of cases studied. This could be the reflection of environmental contamination or it may also be that multiple pathogens act synergistically to produce diarrhea. This result suggests that the future research into

the questions of quantitative association of microorganisms and diarrhoeal disease; synergism between pathogens and serological response to infections may be required to establish the etiology for some pathogens.

ACKNOWLEDGEMENTS

The author would like to thanks Ms. Indu Lamsal, Ms. Sarala Sherchand, Ms Lila Pradhan, Mr Govinda Gurung, Mr Prajjwal Tamrakar and Ms Kamala Lama for their significant help during field and laboratory investigation.

REFERENCE:

- Bern, C., Martinez, J., deZoysa, I. and Glass, R.I. 1992. The magnitude of the global problem of diarrheal disease: a ten-year update. *Bull WHO*. **70**:705-714.
- Black, R.E., Nelson, K.E., Williams, C.M. and Graham, N.M.H. 2001. Diarrheal diseases. *Infectious Disease Epidemiology*. **17**: 497-517.
- Kapikian, A.Z., Kim, H.W., Wyatt, R.G., Cline, W.L., Arrobio, J.O., Brandt, C.D., Rodriquez, W.J., Sack, D.A., Chanock, R.M. and Parrott, R.H. 1976. Human reovirus-like agent as the major pathogen associated with 'winter' gastroenteritis in hospitalized infants and young children. *New Engl. J. Med.* **294**:965-972.
- Kapikian, Z. and Chanock, R.M. 1990 Rotaviruses. In: Virology (Fields BN, Knipe DM, Chanock RM, Hirsch MS, Melnick JL, Monath TP and Roizman B, Eds.). Raven Press, New York. Pp. 1353-1404.
- Greengough, W.B. III. 1982. Bacterial diarrhoeal disease: Current concepts on etiology and pathogenesis. *Southeast Asian J. Trop. Med. Public Health*. **13**: 319-324.
- Lam, S.K. 1982. Current concepts on etiology and pathogenesis of diarrhea caused by viruses. *Southeast Asian J.Trop. Med. Public Health*. **13**: 325-330.
- San Joaquin, V.H. and Marks, M.I. 1982. New agents in diarrhea. *Pediatr. Infect. Dis.* **1**: 53-65.
- Annapurna, E. and Sanyal, S.C. 1977. Enterotoxigenicity of *Aeromonas hydrophila*. *J. Med. Microbiol.* **10**: 317-323.
- Ljungh, A., Popoff, M. and Wadstrom, T. 1977. *Aeromonas hydrophila* in acute diarrhoeal disease: detection of enterotoxin and biotyping of strains. *J. Clin. Microbiol.* **6**: 96-100.
- Saraswati, B., Agrawal, R.K. and Sanyal, S.C. 1983. Further studies on the enteropathogenicity of *Plesiomonas shelloides*. *Indian J. Med. Res.* **78**: 12-18.
- Shlim, D.R., Cohen, M.T., Eaton, M., Ramachandran, R., Long, E.G. and Ungar, B.L.P. 1991. An algae-like organism associated with an outbreak of prolonged diarrhea among foreigners in Nepal. *Am J Trop Med Hyg.* **45**: 383-389.
- Ortega, Y.R., Sterling, C.R., Gillman, R.H., Cama, V.A. and Diaz, F. 1993. *Cyclospora species*-a new protozoan pathogen of humans. *N Engl J Med.* **328**(18): 1308-12.
- Hoge, C.W., Echeverria, P., Rajah, R., Jacobs, J., Malthouse, S., Chapman, E., Jimenez, L.M. and Shlim, D.R. 1995. Prevalence of *Cyclospora* species and other enteric pathogens among children less than 5 years of age in Nepal. *J Clin Microbiol.* **33**(11): 3058-60.
- Shlim, D.R., Hoge, C.W., Rajah, R., Scott, R.M., Pandey, P. and Echeverria, P. 1999. Persistent high risk of diarrhea among foreigners in Nepal during the first 2 years of residence. *Clin Infect Dis.* **29**(3): 613-16.
- Sherchand, J.B. and Cross, J.H. 2001. Emerging pathogen *Cyclospora cayetanensis* infection in Nepal. *Southeast Asian J.Trop.Med.Public Health*. **32**: (suppl 2)143-150.
- Sherchand, J.B. and Cross, J.H. 2004. Parasitic Epidemiological studies of *Cyclospora cayetanensis* in Nepal. *Southeast Asian Journal Trop Med Public Health*. **35**: (Suppl 1) 12-19.

17. De Zoysa, I. and Feachem, R.G. 1985. Interventions for the control of diarrhoeal diseases among young children: rotavirus and cholera immunization. *Bull WHO*. **63**: 569-58.
18. Blacklow, N.R. and Greenberg, H.B. 1991. Medical progress: Viral gastroenteritis. *New Engl. J. Med.* **325**: 252-263.
19. Sherchand, J.B., Larsson, S., Rana, B.J., Dixit, H., Bam, D.S., Adhikari, R.K., Sharma, P.R., Shrestha, M.K. and Shrestha, M.P. 1992. On the incidence of Rotavirus and enteric adenovirus diarrhoea in children attending the outpatient department of Kanti Children's Hospital and general practitioners in the Kathmandu area. *J. Nepalese Med. Assoc.* **30**: 149-53.
20. Sherchand, J.B., Shrestha, M.P., Hommel, M. and Ohara, H. 1998. Hospital based study southern Nepal on morbidity of malaria. *Khoj-Bin: Journal of Nepal Health Research Council.* **2**: 18-22.
21. Adkins, H.J., Escamilla, J., Santiago, L.T., Rañoa, C., Echeverria, P. and Cross, J.H. 1987. Two-year survey of etiologic agents of diarrheal disease at San Lazaro Hospital, Manila, Republic of the Philippines. *J. Clin Microbiol.* **25**: 1143-1147.
22. Albert, M.J., Faruque, S.M., Faruque, A.S.G., Neogi, P.K.B., Ansaruzzaman, M., Bhuiyan, N.A., Alam, K. and Akbar, M.S. 1995. Controlled study of *Escherichia coli* diarrheal infections in Bangladeshi children. *J Clin Microbiol.* **33**: 973-977.
23. Casalino, M., Yusuf, M.W., Nicoletti, M., Bazzicalupo, P., Coppo, A., Colonna, B., Cappelli, C., Bianchini, C., Falbo, V., Ahmed, H.J., Omar, K.H., Maxamuud, K.B. and Maimone, F. 1988. A two-year study of enteric infections associated with diarrhoeal diseases in children in urban Somalia. *Trans R Soc Trop Med Hyg.* **82**:637-641.
24. Mikhail, I.A., Fox, E., Haberberger, R.L. Jr, Ahmed, H. and Abbatte, E.A. 1990. Epidemiology of bacterial pathogens associated with infectious diarrhea in Djibouti. *J Clin Microbiol.* **28**:956-961.
25. Shlim, D.R., Cohen, M.J., Eaton, M., Ramachandran, R., Long, E.G. and Ungar, B.L.P. 1991. An algae-like organism associated with an outbreak of prolonged diarrhea among foreigners in Nepal. *Am. J. Trop. Med. Hyg.* **45**: 383-389.
26. Sherchand, J.B. and Haruki, K. 2004. Rotavirus Study among children and animals of rural and urban communities of Nepal. *Journal of Nepal Health Research Council.* **2**: 5-7.
27. Albert, M.J., Faruque, A.S.G., Faruque, S.M., Sack, R.B. and Mahalanabis, D. 1999. Case-Control Study of Enteropathogens Associated with Childhood Diarrhea in Dhaka, Bangladesh. *J.Clin. Microbiol.* **37**(11): 3458-3464.
28. Ballal, M. and Shivananda, P.G. 2002. Rotavirus and enteric pathogens in infantile diarrhea in Manipal, South India. *Indian J Pediatr.* **69**(5): 393-6.
29. Mertens, T.E., Wijenayake, R., Pinto, M.R., Peiris, J.S., Wijesundera, M.D., Eriyagama, N.B., Karunarathne, K.G. and Ranaweera, L.R. 1990. Microbiological agents associated with childhood diarrhoea in the dry zone of Sri Lanka. *Trop Med Parasitol.* **41**(1): 115-20.
30. Saravanan, P., Ananthan, S. and Ananthasubramanian, M. 2004. Rotavirus infection among infants and young children in Chennai, South India. *Indian Journal of Medical Microbiology.* **22**(4): 212-221.
31. Urbina, D., Arzuza, O., Young, G., Parra, E., Castro, R. and Puello, M. 2003. Rotavirus type A and other enteric pathogens in stool samples from children with acute diarrhea on the Colombian northern coast. *International Journal Microbiol.* **6**(1): 27-32.
32. Zheng, B.J., Ma, G.Z., Tam, J.S.L., Lo, S.K.F., Hon, M., Lam, B.C.C. and Young, C.Y. 1991. The effects of maternal antibodies on neonatal rotavirus infection. *Pediatr Infect Dis Journal.* **10**: 865-868.
33. Mata, L., Simhon, A., Urruta, J.J., Kronmal, R.A., Fernandez, R. and Garcia, B. 1983. Epidemiology of rotaviruses in a cohort of 45 Guatemalan Mayan Indian children observed from birth to the age of three years. *J Infect Dis.* **148**: 452-461.
34. Sharma, A.P. 1978. Quality of drinking water of Kathmandu valley. *J Inst Sci Tech.* **1**: 15-22.
35. Young, B. and Briscoe, J. 1987. A case control study on effect of environmental sanitation on diarrhea morbidity in Malawi. *J Epidemiol Commun Health.* **42**: 83-88.
36. Naficy, A.B., Abu-Elyazeed, R., Holmes, J.L., Rao, M.R., Savarino, S.J., Kim, Y., Wierzba, T.F., Peruski, L., Lee, Y.J., Gensch, J.R., Glass, R.I. and Clemens, J.D. 1999. Epidemiology of rotavirus diarrhea in Egyptian children and implications for disease control. *Am. J. Epidemiol.* **150**: 770-777.
37. LaBaron, C.W., Lew, J., Glass, R.I., Weber, J.M. and Ruiz-Palacios, G.M. 1990. Rotavirus Study Group. Annual rotavirus epidemic patterns in North America. *J. Am. Med. Assoc.* **264**: 983-988.
38. Thomas, F.W., Ibrahim, A.M., Remon, E., Shannond, P., Karim, A.K., Patrick, R., Salwa, F.A., Abdel, F., Kahled, Z., Hind, I.S., John, S. and Robert, F. 2006. Clinic-based surveillance for bacterial and rotavirus associated diarrhea in Egyptian children. *Am. J. Trop. Med. Hyg.* **74**(1): 148-153.
39. Sherchand, J.B. and Cross, J.H. 2001. Emerging pathogen *Cyclospora cayetanensis* infection in Nepal. *Southeast Asian Journal Trop Med Public Health.* **32**: (Suppl 2)143-150.

