

# Antibiotic Susceptibility Pattern of Bacterial Uropathogens causing Urinary Tract Infection in Patients attending Provincial Hospital, Janakpurdham, Nepal

Sanjeet Kumar Jha<sup>1\*</sup>, Saroj Kumar Thakur<sup>2</sup>, Lalit Narayan Yadav<sup>3</sup>, Juni Kumari<sup>3</sup>

<sup>1</sup>Department of Internal Medicine, Provincial Hospital, Madhesh Province, Janakpurdham, Nepal

<sup>2</sup>Department of Microbiology, Provincial Hospital, Madhesh Province, Janakpurdham, Nepal

<sup>3</sup>Department of Biochemistry, Provincial Hospital, Madhesh Province, Janakpurdham, Nepal

## ARTICLE INFO

### Article history:

Received: 29 October 2021

Revised: 11 December 2021

Accepted: 29 December 2021

### \*Correspondence:

Dr. Sanjeet Kumar Jha  
Department of Internal  
Medicine, Provincial Hospital,  
Madhesh Province,  
Janakpurdham, Nepal.

### E-mail:

jhasanjeet0@gmail.com

### Citation:

Jha S K, Thakur S K, Yadav L  
N, Kumari J. Antibiotic  
Susceptibility Pattern of  
Bacterial Uropathogens  
causing Urinary Tract  
Infection in Patients attending  
Provincial Hospital,  
Janakpurdham, Nepal. MedS.  
J. Med. Sci. 2021;1(2):46-50.

## ABSTRACT

**INTRODUCTION:** Distribution and susceptibility of UTI-causing pathogen varies according to place and time. Area specific monitoring studies, aimed to gain knowledge about the type of uropathogens and their susceptibility pattern, help clinicians choose the correct empirical treatment and reduce antibiotic resistance as well as treatment time and financial burden. **MATERIALS AND METHODS:** A hospital based cross-sectional study was carried out in provincial hospital, Madhesh province, Janakpurdham, Nepal from May 2021 to September 2021. Urine samples from 450 clinically suspected cases of UTI were collected and tested bacteriologically following standard procedure. Antibiotic susceptibility test was performed by the disk diffusion according to Clinical Laboratory Standard Institute (CLSI) guidelines. **RESULTS:** Of 450 urine samples, 110 (24.4%) showed significant bacterial growth. E. coli 57 (51.82%) was the most common isolated followed by Pseudomonas aeruginosa 18 (16.36%). Most of the isolates were sensitive to Amikacin (AK), Amoxicillin Clavulanic Acid (AMC), Nitrofurantoin (NIT) whereas most of the gram positive and gram-negative isolates were resistant to Cefixime (CFM), Ampicillin (AMP), Cotrimoxazole (COT). **CONCLUSIONS:** This study showed that E. coli isolates were the predominant uropathogens and AMC is the most effective antibiotic. Most isolates were resistant to Cefixime (CFM), Ampicillin (AMP) and Cotrimoxazole (COT). The presence of highly resistive bacterial isolates, to some of the commonly prescribed drugs, limits the antibiotic prescription options. Drug resistance pattern is an ever-evolving process then isolates become resistant to commonly used drugs. So, frequent surveillance studies are conducted to update clinicians on effectiveness of empirical treatment for UTI.

**Keywords:** Antimicrobial susceptibility, Escherichia coli (E. coli), Urinary tract infection

## INTRODUCTION

Urinary Tract Infection (UTI) refers to the presence of microbial pathogens within the urinary tract leading to an inflammatory response in the epithelium [1]. UTI is the commonest bacterial infection encountered by clinicians with estimated annual global incidence of 250 million [2, 3]. The cases of UTI among Nepalese patients attending hospital ranges from 23.1% to 37.4% [4]. UTI is usually classified by infection site- urine (bacteriuria), bladder (cystitis), kidney (pyelonephritis), prostate (prostatitis) and can be asymptomatic bacteria (ASB) or symptomatic that occurs in a normal genitourinary tract with no prior instrumentation is considered as uncomplicated,

whereas complicated infection is diagnosed in normal genitourinary tract that has either structural or functional abnormalities, including instrumentation [5,6]. More than 95% cases of UTI are due to bacterial causes among which in more than 80% Escherichia coli is the cause [7]. Treatment of UTI cases are often started empirically and therapy is based on the antimicrobial resistance pattern [8]. However, a large proportion of antibiotic usages have contributed resistant bacterial infections resulting in an increased prevalence of antibiotic resistance worldwide [9-12]. Resistance rate and pattern to the most commonly prescribed drugs for UTI treatment varies

considerably in different areas worldwide. The estimation of local etiology and susceptibility profile could support the most effective empirical treatment [13]. Therefore, investigating bacterial distribution pattern and their susceptibility is fundamental for care givers and health planners to guide the expected treatment and interventions. Thus, the aim of this study was to determine bacterial distribution and evaluate their antibiotic susceptibility pattern to commonly used antibiotics.

## MATERIALS AND METHODS

### Study design and setting

This cross-sectional study was conducted from June 2021 to October 2021 on UTI cases, attending provincial hospital Janakpur in outpatient department (OPD), Indoor, Intensive care unit (ICU) and Emergency, under Internal Medicine unit. A total of 450 urine samples were sent for culture, and antibiotic sensitivity test was performed. All the positive culture of 110 cases were included in this study. Patients who received antibiotics within last 15 days or less before sample collection were excluded from the study. Also, patients below the age of 15 years were not included.

### Sample and procedures

### Bacterial isolation, identification and Antibiotic Susceptibility testing

Isolation of uropathogens was performed by a surface streak procedure on Blood agar, Mac Conkey agar, CLED (cystine–lactose–electrolyte-deficient agar or medium) agar & Mannitol salt agar (Hi-media Pvt. Ltd. India) using calibrated loops for semi quantitative method and incubated aerobically at 37<sup>o</sup> C for 24 hours, and those cultures which becomes negative at the end of 24hrs incubations were further incubated for 48hrs [9].

A specimen was considered positive for UTI if a single organism was cultured at a concentration of  $\geq 10^5$  cfu/ml. [9]. Antimicrobial susceptibility of isolates was tested for all bacterial uropathogens by the disk diffusion according to Clinical Laboratory

Standards Institute (CLSI) guidelines [14].ocol following Monica Cheesebrough, 2002 [11].

### Statistical analysis and data management


Data were entered and analyzed in Microsoft Excel [version2010, Microsoft corporation, USA. Frequencies and percentages were expressed. Pie chart was used to present culture positive. Discrete variables were compared using the Chi-square test. Statistical significance were considered at value of the  $p < 0.05$ .

### Ethical considerations

An approval and permission for this study was taken from Provincial hospital before commencing this study.

## RESULTS

Total 450 eligible urine samples were processed among which 110(24.4%) were cultured positive (Figure-1) and hence UTI.



**Table1** Gender and age distribution of UTI patients (n=110)

Age group	Male [n (%)]	Female [n (%)]	Total (%)
15-39	20(18.18)	48(43.64)	68(61.82)
40-59	5(4.55)	2(1.82)	7(6.36)
>=60	25(22.73)	10(9.09)	35(31.82)
Total	50(45.45)	60(54.55)	110(100)
Chi-square=18.48, p-value=0.0001			

**Table 2|** Distribution of bacterial uropathogens isolated from UTI suspected patients (n=110)

Isolates	Number	percentage
E. coli	57	51.82
P. aeruginosa	18	16.36
K. pneumonia	9	8.18
K. oxytoca	4	3.64
Serratia	4	3.64
Staph. Aureus	4	3.64
Acinetobacter	3	2.73

**Table 2| Continued...**

Enterococci	2	1.82
Hafnia	2	1.82
Staph. Epidermidis	2	1.82
Citrobacter	1	0.91
Proteus mirabilis	1	0.91
Serratia marcescens	1	0.91
Staph. saprophyticus	1	0.91
Str. Pyogen	1	0.91
<b>Total</b>	<b>110</b>	<b>100</b>

**Table 3|** Sensitivity pattern of bacterial isolates from urine culture (n=110)

ORAGANISM	NUMBER OF STRAINS (%) SENSITIVE TO														
	AK	AMP	AMC	CPM	CFM	CTR	C	CIP	COT	GEN	IPM	LE	NIT	PTZ	DOX
E. coli (57)	49 (86)	15 (26.3)	46 (80.7)	15 (26.3)	7 (12.3)	13 (22.8)	34 (59.6)	33 (57.9)	7 (12.3)	46 (80.7)	36 (63.2)	40 (70.2)	42 (73.7)	31 (54.4)	14 (24.6)
P. aeruginosa (18)	15 (83.3)	9 (50)	17 (94.4)	0 (0)	0 (0)	0 (0)	12 (66.7)	9 (50)	0 (0)	12 (66.7)	9 (50)	6 (33.3)	14 (77.8)	3 (16.7)	4 (22.2)
K. pneumoniae (9)	7 (77.8)	3 (33.3)	9 (100)	0 (0)	0 (0)	2 (22.2)	4 (44.4)	2 (22.2)	2 (22.2)	1 (11.1)	7 (77.8)	3 (33.3)	7 (77.8)	6 (66.7)	0 (0.0)
K. oxytoca (4)	3 (75)	3 (75)	4 (100)	0 (0)	0 (0)	4 (100)	4 (100)	1 (25)	4 (100)	3 (75)	0 (0)	3 (75)	3 (75.0)	1 (25.0)	0 (0.0)
Serratia (4)	0 (0)	4 (100)	3 (75)	4 (100)	0 (0)	0 (0)	4 (100)	4 (100)	4 (100)	0 (0)	0 (0)	1 (25)	4 (100.0)	4 (100)	0 (0.0)
Staph. Aureus (4)	4 (100)	1 (25)	4 (100)	2 (50)	0 (0)	0 (0)	2 (50)	2 (50)	0 (0)	4 (100)	2 (50)	4 (100)	4 (100.0)	4 (100)	2 (50.0)
Acinetobacter (3)	3 (100)	3 (100)	3 (100)	0 (0)	0 (0)	0 (0)	1 (33.3)	0 (0)	0 (0)	2 (66.7)	2 (66.7)	2 (66.7)	3 (100.0)	0 (0)	1 (33.3)
Enterococci (2)	2 (100)	0 (0)	2 (100)	1 (50)	0 (0)	2 (100)	0 (0)	2 (100)	0 (0)	2 (100)	1 (50)	1 (50.0)	2 (100.0)	1 (50.0)	1 (50.0)
Hafnia (2)	2 (100)	1 (50)	2 (100)	1 (50)	0 (0)	1 (50)	0 (0)	2 (100)	0 (0)	2 (100)	2 (100)	2 (100.0)	1 (50.0)	2 (100.0)	1 (50.0)
Staph. Epidermidis (2)	2 (100)	1 (50)	2 (100)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	2 (100)	1 (50)	2 (100.0)	1 (50.0)	2 (100)	1 (50.0)
Citrobacter (1)	1 (100)	1 (100)	1 (100)	0 (0)	0 (0)	0 (0)	1 (100)	0 (0)	0 (0)	0 (0)	1 (100)	1 (100.0)	1 (100.0)	0	0 (0.0)
Proteus mirabilis (1)	0 (0)	0 (0)	1 (100)	1 (100)	0 (0)	1 (100)	1 (100)	1 (100)	0 (0)	1 (100)	0 (0)	1 (100.0)	1 (100.0)	1 (100.0)	0 (0.0)
Serratia marcescens (1)	0 (0)	1 (100)	1 (100)	1 (100)	0 (0)	0 (0)	1 (100)	1 (100)	1 (100)	0 (0)	0 (0)	0 (0.0)	1 (100.0)	1 (100.0)	0 (0.0)
Staph. Saprophyticus (1)	1 (100)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (100)	0 (0)	0 (0)	1 (100)	1 (100)	1 (100.0)	0 (0.0)	1 (100.0)	0 (0.0)
Str. Pyogens (1)	1 (100.0)	1 (100)	1 (100)	0 (0)	0 (0)	0 (0)	1 (100)	0 (0)	0 (0)	1 (100)	0 (0)	1 (100.0)	1 (100.0)	1 (100.0)	0 (0.0)
Total (110)	90 (81.8)	43 (39.1)	96 (87.3)	25 (22.7)	7 (6.4)	23 (20.9)	65 (59.1)	57 (51.8)	18 (16.4)	77 (70)	62 (56.4)	68 (61.8)	85 (77.3)	58 (52.7)	24 (25.8)

Of 110 cases, there were 50(45.5%) samples of male and 60(54.5%) of female. These results indicate that the occurrence of UTI was higher in female patients than in males patients. The highest susceptible to UTI was below 40 years age group of patients (61.82%) followed by 60 or more years (31.82%). The highest occurrence of UTI in female was found below the age of 40 years (43.64%). However, in male the highest susceptible age group to UTI was above 60 years (22.73%). A significant association was observed for gender and age group ( $p=0.0001$ ) among UTI patients [Table1].

**Out of total 110** bacterial isolates in urine samples, 90.91% were Gram negative and 9.09% were Gram positive. *E. coli* was found to be the most dominant bacteria among all isolated uropathogens with the occurrence of 57(51.82%), followed by *Pseudomonas aeruginosa* 18(16.36%) and *Klebsiella pneumoniae* 9 (8.18%). Among Gram positive staphylococcus, aureus was most common uropathogens (3.64%) (Table 2).

**Bacterial uropathogens** isolated from patients with UTI revealed the presence of high level of single and multiple antibiotic sensitivity against commonly prescribed drugs shown in Table 3. *E. coli* which was the most common cause of UTI showed high percentage of sensitivity to Amikacin (AK) (86%) followed by Gentamycin (GEN) (80.7%), Amoxicillin clavulanic acid (AMC) (80.7%), and Nitrofurantoin (NIT) (73.7%) whereas cefixime (CFM) and cotrimoxazole (COT) were least sensitive (12.3%).

## DISCUSSION

Bacterial infection of the urinary tract is one of the common morbidities seeking medical attention [15]. Increasing antimicrobial resistance has been documented worldwide [16-20]. This study provides valuable data to compare and monitor the status of antibiotic sensitivity among uropathogens to improve empirical treatment. In this study, the isolation rate of bacterial from urine was 24.44% which is similar to other studies that accounts for 25.6% [21], 22% [22], though it is lesser than 38.6% [23] and 35.5% [24].

## ADDITIONAL INFORMATION AND DECLARATIONS

**Acknowledgements:** Authors wish to thank medical superintendent of Provincial hospital for all their supports during this study.

**Competing Interests:** The authors declare no competing interests.

Our study showed a higher occurrence of UTI in females (54.5%) than in males (45.5%) which correlates with other findings [25-28]. The reason behind high prevalence in female is due to close proximity of urethral meatus to the anus, short urethra. *E. coli* (51.82%) is the most common bacteria isolated from urine samples and this finding is in agreement with others finding too [13, 29, 30]. Second most common isolated bacterium was *pseudomonas aeruginosa* (16.36%) which was in agreement with some other studies [31, 32].

Though it does not correlate with some studies [33] in which *Klebsiella* species was the second most common. Resistance to antibiotic has been noted since the first use of these agents and is a worldwide increasing problem [17-20]. *E. coli*, *Pseudomonas* are highly sensitive to Amikacin (AK), Amoxicillin clavulanic acid (AMC) and Nitrofurantoin (NIT).

A comparable rate of sensitivity has been reported to these drugs in some previous studies [29,33]. Thus, these drugs could be considered as alternative options in the empirical treatment of UTIs. The cefixime (CFM), Ampicillin (AMP) were found resistant to most of the antimicrobials. So, these drugs should not be used as the empirical treatment for UTI in this community.

**CONCLUSIONS** *E. coli* was the most common isolated organism in UTI patients. Bacterial uropathogens isolated from patients with UTI revealed the presence of high level of single and multiple antibiotic sensitivity against commonly prescribed drugs. Most of the isolates were sensitive to Amikacin, Amoxycillin Clavulanic Acid, Nitrofurantoin whereas most of the gram positive and gram-negative isolates were resistant to Cefixime, Ampicillin, Cotrimoxazole. The isolation of bacterial pathogens with sensitivity pattern of antimicrobial provides valuable data to improve recommendation in specific community for empirical treatment of UTI. As drug sensitivity and resistance is an evolving process, routine surveillance studies should be conducted to guide clinicians with knowledge about the most effective empirical treatment of UTIs.

**Funding:** Self-funded

**Author Contributions:** Concept and design: S.K.J, Statistical analysis: S.K.J, Writing of the manuscript: L.N.Y, J.K. Data collection: L.N.Y, S.K.T and J.K. Revision and editing: S.K.J. All

authors have read and agreed with the contents of the final manuscript towards publication.

**Data Availability:** Data will be available upon request to corresponding authors after valid reason.

## REFERENCES

- Nature, Macmillan Publisher Limited. Urinary tract infection. Available at: <http://www.nature.com/urinary-tract-infection>.
- Ronald AR, Nicolle LE, Stam E, et al. Urinary tract infection in adults: research priorities and strategies. *Int J Antimicrob Agents*. 2001; 17:343-348.
- Barišić Z, Babić-Erceg A, Brozić El, et al. Urinary tract infection in South Croatia: Aetiology and antimicrobial. *Intl J Antimicrob Agents*. 2003; 22: s61-s64.
- Rai GK, Upreti HC, Rai SK, Shah KP, Shrestha RM. Causative agents of urinary tract infections in children and their antibiotic sensitivity pattern: a hospital passed study. *Nepal Med Coll J*. 2008; 10(2):86-90.
- Gonzalez CM, Schaeffer AJ. Treatment of urinary tract infection: what's old, what's new, and what works. *World J Urol*. 1999; 17: 372-382.
- Stamm WE, Hooton TM. Management of urinary tract infections in adult. *N Engl J Med*. 1993; 329:1328-1334.
- Nachimuthu R, Chettipalayam S, Velramar B, Kurumandur R, Velu R. Urinary tract infection and Antimicrobial susceptibility pattern of extended spectrum beta lactamase producing clinical isolates. *Adv Biol Res*. 2008; 2(5-6): 78-82.
- Wilson ML, Gaido L. Laboratory Diagnosis of Urinary Tract Infection in Adult Patients. *Clin Infect Dis* 2004; 38: 1150-1158.
- Bonadio M, Meini M, Spetaleri P, Gilgi C. Current microbiological and clinical aspects of urinary tract infections. *Eur J Urol*. 2001; 40:439-445.
- National committee for clinical laboratory standards. Performance standards for antimicrobial disc susceptibility test. 7thed. Wayne Pennsylvania, USA: NCCLS; 2000. M2-A7.
- Grude N, Tveten Y, Kristiansen BE. Urinary tract infections in Norway: bacterial etiology and susceptibility, a retrospective study of clinical isolates. *Clin Microbial Infect*. 2001; 7: 543-547.
- Kripke C. Duration of therapy for women with uncomplicated UTI. *Am Fam Physician*. 2005; 72: 2219.
- Farajnia S, Alikhani MY, Ghotaslou R, Naghili B, Nakhband A. Causative agents and antimicrobial susceptibilities of urinary tract infection in the Northwest of Iran. *Int J Infect Dis*. 2009; 13:140-144.
- Clinal and Laboratory Standard Institute. Supplemental table. Performance standards for antimicrobial susceptibility testing; 15th informational supplement. CLSI publication. M100-S15, M2 -A8, M7-A6 Pennsylvania: CLSI;2005.
- Kebira AN, Ochola P, and Khanadi SA. Isolation and antimicrobial susceptibility testing of Escherichia coli causing urinary tract infections. *J Appl Biosci*. 2009; 1320-1325.
- Jani M, Shah S, And Prajapati S. Antibacterial screening and qualitative phytochemical estimation of selected aquatic plants. *Advances in biological research*. 2012; 6 (1) 19-23.
- Kashef N, David GE, Shahbazi S. Antimicrobial susceptibility patterns of community acquired uropathogens in Tehran, Iran . *J Infect in developing country*. 2010; 4(1): 202-206.
- Karlowsky JA, Jones ME, Thornsberry C, Critchley I, Kelly LJ, Sham DF. Prevalence of antimicrobial resistance among urinary tract pathogens isolated from female out patients across the USA in1999. *Int J Antimicrobial Agents*. 2001; 18(20): 121-127.
- Rajalakshmi V, Amsaveni V. Antibiotic susceptibility of bacterial pathogens isolated from diabetic patients. *Int J Microbio Res*. 2012; 3(1) 30-32.
- Rashedmarandi FRM, Saremi M. A survey on urinary pathogens and their antimicrobial susceptibility among patient with significant bacteriuria. *Iranian J path*. 2008; 3: 191-196.
- Nedolisa. Bacteriology of urinary tract infection amongst patients attending JO University Teaching Hospital . JUTH. 1998 [M. S Thesis].
- Ekweonzor CC, Onyemenen TN. UTI in Ibadan, Nigeria : causative organism and antimicrobial sensitivity pattern. *African J Med Sci*. 1996; 25: 165-169.
- Akinyemi O, Alabi TN, Taiwo MA, and Omonigbehin EA. Antimicrobial susceptibility pattern and plasmid profiles of pathogenic bacteria isolated from subjects with urinary tract infection in Lagos, Nigeria. *Nigerian quat J of Hospital Med*. 1997; 1: 7-11.
- Ebie MY, Kandaki-Olukemi YT, Ayanbadejo J, and Tanyigna KB. UTI infections in Nigerian Military Hospital. *Nigerian J microbiology*. 2001; 15:31-37.
- Schaeffer AJ, Rajan N, Cao Q, et al. Host pathogenesis in urinary tract infections. *Int J Antimicrobial Agents*. 2001; 17(4): 245-251.
- Orrett FA. Urinary tract infections in general practice in rural community in South Trinidad. *Saudi Med J*. 2001; 22(6): 537-540.
- Boucher HW, Talbot GH, Bradley JS et al. Bad bugs, no drugs: no ESKAPE! An update from infectious diseases society of America. *Clinical Infectious Disease*. 2009; 48(1): 1-12
- Oladeinde BH, Omaregie R, Olley M and Anunibe JA. Urinary tract infection in a rural community of Nigeria. *North American journal of medical sciences*. 2011; 3(2): 75-77.
- Tessema B, Kassu A, Mulu A, Yismaw G. Predominant isolates of urinary tract pathogens and their susceptibility pattern in Gonder University Teaching Hospital, Northwest Ethiopia. *Ethiop Med J*. 2007; 45:61-67.
- Dromigny JA, Nabeth P, Perrier Gros Claude JD. Distribution and susceptibility of bacterial urinary tract infection in Dakar, Senegal. *Int J Antimicrob Agents*. 2002; 20: 339-347
- TambekarDH, Dhanorkar DV, Gulhane SR, Khandelwal VK, and Dudhane MN. Antibacterial susceptibility of some urinary tract pathogens to commonly used antibiotics. *African journal of Biotechnology*. 2006; 5(17): 1562-1565.
- Kolawole AS, Kolawole AM, Kandaki-Olukemi YT, Babatunde SK, Durowade KA, and Kolawole CF. prevalence of urinary tract infection (UTI) among patients attending Dalhatu Araf Specialist Hospital, Lafia, Nigeria. *Int J Medicinal Med Sci*. 2009; 1(5): 163-167.
- Moges F, Genetu A. Antibiotic sensitivity of common bacteria pathogens in urinary tract infection at Gonder Hospital, Ethiopia. *Ethiop J Health Dev*. 2009; 79: 140-1

### Publisher's Note

MJMMS remains neutral with regard to jurisdictional claims in published materials and institutional affiliations.



CCREACH will help you at every step for the manuscript submitted to MJMMS.

- We accept pre-submission inquiries.
- We provide round the clock customer support
- Convenient online submission
- Plagiarism check
- Rigorous peer review
- Indexed in NepJOL and other indexing services
- Maximum visibility for your research
- Open access

Submit your manuscript at:  
Website: [www.medsjournal.org](http://www.medsjournal.org)  
e-mail: [editormjms@gmail.com](mailto:editormjms@gmail.com)

