

Effect of Different Postharvest Treatments on Prolonging Shelf life of *Citrus reticulata* Blanco.

Pabitra Joshi¹, Bishnu Raj Ojha¹ and Arun Kafle²

¹Agriculture and Forestry University, Rampur, Chitwan,

²Vegetable Crops Development Centre, Khumaltar, Lalitpur

Corresponding author's email address: pabitrajoshi77@gmail.com

Received on: 16 July 2019, Revised on: 10 November 2019, Accepted on: 5 December 2019

Abstract

This study was conducted during January to April 2018 to evaluate the effect of different post-harvest treatments on maintaining quality and shelf life of mandarin. Laboratory experiment was conducted under Complete Randomized Design (CRD) with four replications and seven treatments (T1 = Control i.e. dipped in distilled water, T2 = Cinnamon oil @2%, T3 = Eucalyptus oil @2%, T4 = Calcium chloride @1%, T5 = Bavistin @0.1%, T6 = Paraffin Wax @10%, T7 = Paraffin Wax @ 10% + Bavistin @0.1%). Data were recorded in every 2 days interval and the final data was taken in 13 days interval because of the limited destructive sample. Postharvest treatment with wax @10% in combination with Bavistin @0.1% had minimum physiological loss in weight (6.61%) and maximum juice recovery percentage (43.72%) which was statistically at par with wax (10%), Bavistin (0.1%), Cinnamon oil (2%), Eucalyptus oil (2%) and Calcium chloride (1%) treated fruits. Also at the end of storage period the highest Total soluble solute (TSS) content (15.45°Brix) was recorded in Calcium chloride (1%) treated fruits which were statistically at par with control, Eucalyptus oil (2%), Bavistin (0.1%) treated fruits. The pH was found non-significant throughout the storage period whereas postharvest life was found the maximum (73 Days) in fruits treated with wax (10%) in combination with Bavistin (0.1%) while it was only 46 days in control.

Keywords: Acidity, Ascorbic acid, Carotenoids, Lycopene, Dehydration ratio

Introduction

Mandarin (*Citrus reticulata* Blanco.) is a group name for a class of oranges belonging to family Rutaceae with bright colored peel and pulp, excellent flavor, easy-to-peel rind and segments that separate easily, is believed to have originated in Southeastern Asia (Parashar, 2009). In Nepal, the total area production and productivity of mandarin orange is 16,248 hectare, 146,690 Metric tons and 9.0Mt/ha which is very low compared to the most citrus growing countries in the world (MoAD, 2016). Low productivity, decline in quality and heavy post-harvest losses are key hurdles faced by Nepalese fruit industry (Rokaya, 2017).

Nearly 20-25 percent of mandarin fruits are wasted due to faulty postharvest management i.e. 7% during harvesting, 25% during transportation, 3% while grading, 10% in packaging, and 5% during marketing (Bhattarai et al. 2013). Since fruits are perishable in nature they cannot be kept for a long-time during transportation and storage. Due to low adaptation of improved techniques during pre and post-harvest stage, both external and internal chemical quality attributes are lost. The postharvest losses can be minimized by extending shelf life through checking the rate of transpiration, respiration, microbial infection & protecting membranes from disorganization (Sahu, 2016). Among the different methods used to extend the shelf life alternative of low-cost technology i.e. the

application of the edible coating (oil, wax, chemical) to fruit has received attention worldwide as these coatings are maintaining quality even under ordinary storage condition (Bisen et al. 2012). Edible coating of fruits can result in the creation of a modified atmosphere due to blockage of the pores within the fruits, reducing respiration rate and improving postharvest quality (Kader, 2005). Considering the above facts this study was carried out with the general objective of evaluating the effect of different postharvest treatments on fruit quality and shelf life of mandarin under room condition.

Methodology

Mandarin fruit of Khoku Local cultivar (local mandarin of Dhankuta) with uniform size, healthy, greenish yellow and well matured from the private orchard of Udayapur district (Katari Municipality, Katunje, one of the potential districts for mandarin cultivation in Nepal) were selected and harvested by clipper keeping with small pedicel intact and collected in crates from the orchard on Poush 25, 2074 (January 09, 2018). The collected fruits were sorted and graded on the basis of size (uniform medium sized) and maturity (greenish yellow in color) for treatments. The fruits were stored under an ambient condition at Beteni lab (PMAMP Citrus Zone) which is located 1300 masl at Udayapur District, Nepal. The experiment was conducted from January to April, 2018 which was laid out in CRD with 7 treatments each replicated 4 times. The treatments applied were T1: Control (distilled water), T2: Cinnamon oil@2%, T3: Eucalyptus oil @2%, T4: Calcium Chloride (CaCl₂)@1%, T5: Bavistin @0.1%, T6: Wax emulsion @10%, T7: Wax (10%) in combination with Bavistin (0.1%). Different Essential oils (Cinnamon oil, Eucalyptus oil) at 2 % concentration was prepared. The fungicidal solution of Bavistin

Titration acidity (TA)

The acidity of the fruits from each treatment was estimated as per standard procedures of AOAC (2005). A total of 10 ml of the clear juice of a fruit from each

$$\text{Titration Acidity (\%)} = \frac{\text{ml of NaOH used} \times \text{acid factor (i.e 0.0064 for citric acid)} \times 100}{\text{Volume (ml) of Juice used}}$$

TSS/TA

TSS /TA ratio was calculated by dividing the TSS

$$\text{TSS /TA} = \frac{\text{Total soluble solids}}{\text{Titration acidity}}$$

@ 0.1 % was prepared by dissolving 1 g of Bavistin (amorphous) 1000 ml of distilled water. This emulsifier wax solution was prepared as a procedure outlined by (Rokaya, 2017). Fruits from each treatment separately dipped for 2 minutes in each prepared solution in a bucket and were dried for 5 minutes under the shed over the newspaper. After the treatment completion, 10 fruits from each treatment were allocated in the plastic tray as a destructive sample from which every time one fruit per replication was used for analysis and 5 fruits of each treatment were allocated in each tray as the non-destructive sample under the ambient condition 12.42±0.280C mean temperature, 68.56±1.46% RH for 41 days where temperature and RH was measured with the help of digital thermo-hygrometer (ERMA) three times a day and was averaged for the whole duration. Observations were recorded in 2 days interval up to 27 days and final observation was taken at 41 day (because of limited sample). Following parameters were evaluated during the storage period.

Physiological Loss in weight (%)

Weight loss was recorded in the same 5 fruits (non-destructive sample). A digital sensitive balance was used to determine fruit weight. The weight loss was calculated according to the formula:

$$W1 = [(W0 - Wt) / W0] \times 100 \%$$

Where W1 is the percentage weight loss,

W0 is the weight of the initial fruit and

Wt is the weight of the fruits at the designated time.

Total Soluble Solid (0Brix)

Total soluble solids (0Brix) was determined with the help of Erma hand-held refractometer.

treatment was taken and titrated against standard 0.1 N of sodium hydroxide (NaOH) solution using phenolphthalein as an indicator. Then the titration acidity of the fruit was expressed in percentage using the following formula:

content by titration acidity of each treatment and average was recorded. Following formula was used to calculate TSS/TA ratio:

pH of the juice

pH of the juice was measured with the help of digital pH meter.

Temperature & RH

Temperature and RH were recorded each day during the experimental period using ERMA thermo-hygrometer.

Juice content

Juice was extracted by squeezing by hands. The volume of juice was measured (ml/fruit) using beaker. Average juice percentage per fruit was obtained from the following formula:

$$\text{Juice (\%)} = \frac{\text{Juice weight per fruit} \times 100}{\text{Individual fruit weight}}$$

Statistical method

The data pertaining to various parameters were collected at different stages and intervals and tabulated in an Excel sheet for analysis as mentioned by Gomez and Gomez (1984). All routine statistical analysis was carried out using Genstat software 15th Edition. This software was used to generate (LSD) test at 0.05 ($p < 0.05$) by Analysis of Variance (ANOVA) to determine the significant difference among the treatment means.

Results and Discussions

Physiological loss in weight (PLW)

The physiological loss in weight (PLW) was significantly increased in all the treatments with the advancement of the storage period and the increasing trend in the weight loss percentage was found the maximum in control up to 41 days of storage. Minimum percentage of PLW was observed in the fruits treated with wax in combination with Bavistin during the whole storage period and the losses ranged from 0.90% in 6th day to 6.61% in 41th day whereas maximum weight loss was recorded in the fruits with untreated as control (1.59% to 18.09%) which was statistically at par with the findings of Calcium chloride (1%), Eucalyptus oil (2%) and Bavistin (0.1%) during the storage (Table 1).

This minimum weight loss in the wax-treated fruits was might be due to retardation in the process of transpiration and respiration by the closing of lenticels and stomata of the cell wall of the fruit skin. Thus wax emulsion might have been an effective treatment to reduce weight loss. Wax coated fruits retained better glossiness and fresh appearance being a moisture barrier in the study carried by (Mahajan et al. 2005). A study in tangerine citrus var. Siam Banjar showed that the application of wax coatings in combination with low-temperature storage proved effective in reducing the percentage weight loss (Hassan et al. 2014). The findings were in consonance with the findings of the Ahmad et al. (2013) in Kinnow fruit, Rokaya (2017) in Mandarin, Sahu (2016) in custard apple, who found minimum weight loss in the fruits treated with a wax emulsion.

Table 1: Effect of postharvest treatments on physiological weight loss (%) of mandarin fruit during storage at ambient condition ($12.42 \pm 0.280^\circ\text{C}$ mean temperature, $68.56 \pm 1.46\%$ RH), Udayapur, Nepal, (2017-18)

Treatments	The physiological loss in weight (%) on days indicated									
	6DAY	9DAY	12DAY	15DAY	18DAY	21DAY	24DAY	27DAY	30DAY	41DAY
Control	1.59abc	3.81a	6.03a	6.98a	8.25a	9.21a	10.79a	12.06a	14.29ab	18.09a
Cinnamon Oil (2%)	1.25bcd	2.19b	4.70bc	5.95ab	6.89ab	7.52ab	9.09ab	9.71bc	10.28c	12.70b
Eucalyptus oil (2%)	1.56abc	2.81ab	4.06c	5.01b	5.94b	6.88b	7.82b	9.07c	12.20bc	14.38ab
CaCl ₂ (1%)	2.17ab	3.41a	6.19a	6.81a	7.74a	8.67ab	10.22a	11.15ab	15.17a	17.03a
Bavistin (0.1%)	2.23a	3.51a	5.73ab	6.05ab	7.32ab	8.28ab	10.51a	11.47ab	14.97a	17.83a
Wax (10%)	0.60d	0.90c	2.09d	2.09c	2.40c	2.99c	3.89c	4.19d	6.58d	11.98b
Wax (10%) + Bavistin (0.1%)	0.90cd	1.20c	1.80d	2.10c	3.01c	3.01c	3.91c	3.91d	5.41d	6.61c
SEm (\pm)	0.293	0.293	0.326	0.354	0.532	0.458	0.592	0.539	0.759	1.227
LSD (=0.05)	0.862	0.959	1.042	1.564	1.347	1.743	1.584	1.721	2.232	3.608
CV, %	39.8	25.6	16.2	21.3	15.4	17.8	13.4	13.3	13.5	17.4
P-value	0.005	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Grand mean	1.47	2.55	4.37	5.00	5.94	6.65	8.03	8.79	11.27	14.09

LSD = Least Significant Difference, SEM = Standard Error of Mean, and CV = Coefficient of variation, ns = non-significant

Juice content

Table 2 shows that the juice recovery percentage was decreased with time during the storage in all the treatments but was not significantly lower. Wax + Bavistin treated fruits recorded the maximum juice recovery percentage (43.72%) which was statistically at par with the findings of Wax 10%, Bavistin 0.1%, Eucalyptus oil (2%), Cinnamon oil (2%) and Calcium chloride 1% at 4th day of storage whereas the minimum juice recovery percentage (33.89%) was observed in control fruits.

The trend of decrease in juice percentage during the storage was might be due to loss of moisture from the surface of the fruits. The wax treated fruits in combination with Bavistin showed a low reduction in juice content during storage as compared to control or other essential oils. This might be due to the fact that the wax acted as a barrier which had checked the losses of the moisture from the fruit surface. Ahamad

et al. (2013) reported higher juice recovery percentage in PE-packed (Polyethylene packed) fruits (T10) followed by the fruits with 100% Sta-Fresh 960 (T4) (Trade name of a commercial edible coating material) which might be due to less water loss in PE-packaging and waxing treatments as the combination acts as a barrier to moisture loss. The minimum decrease in juice percentage was observed in the fruits treated with wax 10% plus Bavistin 0.1% from the 1st week (49.56%) to the 4th week (43.81%) followed by wax 10% from the 1st week (49.49%) to the 4th week (43.45%) as against control from the 1st week (47.26%) to the 4th week (34.65%). Rokaya (2017) reported that the fruits treated with wax and in combination with Bavistin showed low reduction in juice content during storage as compared to other chemically treated fruits and control which might be due the fact that the wax acted as a barrier which had checked the losses of the moisture from the fruit surface. These results are in line with Mahajan et al. (2013) in Kinnow fruit, Bisen et al. (2012) in Kagzi lime.

Table 2: Effect of postharvest treatments on juice recovery (%) of mandarin fruit during storage at ambient condition ($12.42 \pm 0.280^\circ\text{C}$ mean temperature, $68.56 \pm 1.46\%$ RH), Udayapur, Nepal, (2017-18)

Treatments	Juice recovery % of fruit on days indicated									
	Day 3	Day 6	Day 9	Day 12	Day 15	Day 18	Day 21	Day 24	Day 27	Day 41
Control	46.60	45.89	44.61	44.26	43.72	42.27	41.02	39.82	38.00	33.89b
Cinnamon Oil (2%)	46.81	46.47	46.33	44.60	44.56	43.83	42.43	41.52	40.61	37.07ab
Eucalyptus oil (2%)	48.23	47.67	47.64	46.95	45.70	45.50	44.81	44.80	38.62	37.80ab
CaCl ₂ (1%)	45.61	45.57	45.12	44.97	44.41	43.87	43.30	42.68	41.93	41.47a
Bavistin (0.1%)	46.69	45.65	45.23	44.56	44.49	43.62	42.72	42.68	42.45	41.86a
Wax (10%)	48.22	47.79	47.09	45.09	45.34	44.33	43.30	43.16	42.89	42.04a
Wax (10%) + Bavistin (0.1%)	48.23	47.07	46.37	46.05	45.62	45.00	44.66	44.35	44.27	43.72a
SEm (\pm)	1.731	1.933	1.855	1.282	2.044	1.133	1.565	1.807	1.869	2.173
LSD (=0.05)	ns	ns	ns	ns	ns	Ns	ns	ns	ns	6.390
CV, %	7.3	8.3	8.1	5.7	9.1	5.1	7.2	8.5	9.1	10.9
P-value	0.896	0.961	0.897	0.759	0.991	0.550	0.653	0.534	0.225	0.048
Grand mean	47.20	46.59	46.06	45.21	44.83	44.06	43.18	42.71	41.25	39.69

LSD = Least Significant Difference, SEM = Standard Error of Mean, and CV = Coefficient of variation, ns = non-significant

Total soluble solutes (TSS)

TSS content is one of the major indicators that determine the quality of mandarin orange. As shown in Table 3, TSS content increased with the increasing period of

storage in all the treatments and the increasing trend is higher in untreated (control), Eucalyptus and calcium chloride treated fruits than the fruits treated with other coatings. Fruits treated with calcium chloride showed the maximum TSS content during the storage

period and ranged from 12.30°brix during 3rd day to 15.45°brix during 41st day which was statistically at par with untreated fruits & Eucalyptus treated fruits and minimum TSS content was recorded in the fruits treated with 10% wax from 3rd day of storage (11.83°brix) to 41st day of storage (13.35°brix) which was statistically at par with wax 10% in combination with Bavistin 0.1% from 3rd day to 41st day of storage. The trend showed that wax treated fruits was significantly superior because of the gradual increment in the TSS change whereas in calcium chloride treated and control it was increased at a faster pace. The faster rate in the TSS increment in the calcium chloride treated and untreated fruits were might be due to faster metabolic activities through respiration and transpiration than in the other

coatings (Akhtar et al., 2010). (Rokaya, 2017) reported that untreated (control) fruits showed the maximum TSS content during the storage and ranged from the 1st week (10.92° Brix) to the 4th week (12.88° Brix) and minimum TSS was recorded in the fruits treated with wax 10% from the 1st week (10.35° Brix) to the 4th week (11.51° Brix) which was at par with wax 10% in combination with Bavistin 0.1% from the 1st week (10.39° Brix) to the 4th week (11.65° Brix). Similarly, the results are in line with the results of Bisen et al. (2012) in Kagzi lime; Waskar & Gaikwad (2005) in Keshar mango; Jhologiker & Reddy (2007) in Sugar apple (*Annona squamosa* L.) fruits, Shahid & Abbasi (2011) in sweet orange, Hassan et al. (2014) in tangerine citrus.

Table 3: Effect of postharvest treatments on total soluble solids (TSS) of mandarin fruit during storage at ambient condition (12.42±0.280C mean temperature, 68.56±1.46% RH), Udayapur, Nepal, (2017-18)

Treatments	TSS of fruits on days indicated									
	Day 3	Day 6	Day 9	Day 12	Day 15	Day 18	Day 21	Day 24	Day 27	Day 41
Control	11.85	12.23	12.55	13.00	13.18	13.70	13.93	14.23 ^a	14.25 ^a	15.35 ^a
Cinnamon Oil (2%)	11.80	11.83	12.60	12.75	12.85	13.50	13.53	13.60 ^{abc}	13.63 ^{ab}	14.05 ^{bc}
Eucalyptus oil (2%)	12.50	12.65	12.90	13.03	13.08	13.50	13.62	13.63 ^{abc}	13.65 ^{ab}	14.53 ^{ab}
CaCl ₂ (1%)	12.30	12.38	13.28	13.30	13.33	13.60	13.68	14.03 ^a	14.13 ^a	15.45 ^a
Bavistin (0.1%)	12.00	12.20	12.63	12.73	12.88	13.70	13.73	13.90 ^{ab}	13.95 ^a	14.60 ^{ab}
Wax (10%)	11.83	11.95	12.10	12.30	12.40	12.65	12.78	13.03 ^{bc}	13.10 ^b	13.35 ^c
Wax (10%) + Bavistin (0.1%)	11.93	12.00	12.33	12.50	12.68	12.83	12.90	13.00 ^c	13.08 ^b	13.12 ^c
Sem (±)	0.663	0.389	0.370	0.318	0.286	0.390	0.416	0.276	0.251	0.347
LSD (=0.05)	ns	ns	ns	ns	ns	Ns	ns	0.811	0.738	1.021
CV, %	11.0	6.4	5.9	5.0	4.4	5.8	6.2	4.0	3.7	4.8
P-value	0.984	0.787	0.417	0.376	0.342	0.332	0.394	0.030	0.016	<0.001
Grand mean	12.03	12.18	12.63	12.80	12.91	13.35	13.45	13.63	13.68	14.35

LSD = Least Significant Difference, SEM = Standard Error of Mean, and CV = Coefficient of variation, ns = non-significant

Titrate Acidity

As data presented in Table 4 titratable acidity of different treatments was significant at the end of the storage period. The TA was significantly decreased with the advancement of the storage period. Higher acidity was recorded in the fruits treated with wax 10% along with Bavistin 0.1% which was 0.68 at 3rd day of storage and 0.55 on 41th day of storage period whereas there was a significant decrease in the TA content of the fruits left untreated which was 0.66 on 3rd day of storage and 0.40 on 41st day of storage period. The decreasing trend of titratable acidity during the storage

period was probably due to the utilization of acid in the tricarboxylic acid cycle in the respiration process (Rokaya, 2017). The higher acidity in wax treated fruits was might be due to less utilization of the acids in the respiration process during the storage whereas untreated fruits with minimum acids was due to faster utilization of the acids in the respiration process during storage (Bisen et al. 2012). Rokaya (2017) recorded maximum TA in the fruits treated with wax 10% plus Bavistin 0.1% against control at the end of the storage. Similarly, the results are in line with, Ahmad et al. (2013) in Kinnow fruit, Hassan et al. (2014) tangerine citrus var. Siam Banjar.

Table 4: Effect of postharvest treatments on titratable acidity (TA) of mandarin fruit during storage at ambient condition ($12.42 \pm 0.280^\circ\text{C}$ mean temperature, $68.56 \pm 1.46\%$ RH), Udayapur, Nepal, (2017-18)

Treatments	Titrable acidity on days indicated (%)									
	Day 3	Day 6	Day 9	Day 12	Day 15	Day 18	Day 21	Day 24	Day 27	Day 41
Control	0.66	0.63	0.61	0.60	0.57	0.54	0.50	0.47	0.42b	0.40c
Cinnamon Oil (2%)	0.68	0.67	0.65	0.64	0.62	0.60	0.59	0.58	0.51a	0.48abc
Eucalyptus oil (2%)	0.68	0.67	0.63	0.62	0.60	0.58	0.56	0.54	0.53a	0.50abc
CaCl ₂ (1%)	0.67	0.64	0.61	0.58	0.56	0.55	0.52	0.50	0.49ab	0.47abc
Bavistin (0.1%)	0.67	0.66	0.64	0.58	0.56	0.54	0.51	0.49	0.46ab	0.43bc
Wax (10%)	0.69	0.66	0.64	0.62	0.60	0.59	0.56	0.55	0.54a	0.52ab
Wax (10%) + Bavistin (0.1%)	0.68	0.66	0.64	0.63	0.62	0.60	0.58	0.57	0.56a	0.55a
Sem (\pm)	0.0303	0.0543	0.0431	0.0280	0.0382	0.0357	0.0653	0.0343	0.0290	0.0330
LSD (=0.05)	ns	Ns	ns	ns	ns	Ns	ns	ns	0.0852	0.0970
CV, %	9.0	16.6	13.6	9.2	13.0	12.5	24.0	13.0	11.5	13.8
P-value	0.989	0.998	0.989	0.683	0.827	0.724	0.929	0.199	0.030	0.067
Grand mean	0.68	0.66	0.63	0.61	0.59	0.57	0.54	0.53	0.50	0.48

LSD = Least Significant Difference, SEM = Standard Error of Mean, and CV = Coefficient of variation, ns = non-significant

TSS/TA Ratio

TSS/TA ratio of mandarin fruit as influenced by a different combination of treatments is depicted in Table 5. At the beginning of the storage period from 3rd day to 21st day no significant differences were observed among the treatments. From 24th day to

41st day of storage significant differences among the treatments were observed. On the 41st day of storage, the significantly maximum ratio was observed with untreated fruits (38.06) while the minimum ratio was observed with wax 10% in combination with Bavistin 0.1% (24.73).

Table 5: Effect of postharvest treatments on TSS/TA ratio of mandarin fruit during storage at ambient condition ($12.42 \pm 0.280^\circ\text{C}$ mean temperature, $68.56 \pm 1.46\%$ RH), Udayapur, Nepal, (2017-18)

Treatments	TSS/TA on days indicated									
	Day 3	Day 6	Day 9	Day 12	Day 15	Day 18	Day 21	Day 24	Day 27	Day 41
Control	17.78	19.44	20.68	22.00	23.46	25.72	28.36	30.68a	34.35a	38.06a
Cinnamon Oil (2%)	17.37	18.13	19.66	19.93	20.88	22.96	23.08	23.38b	27.26bc	30.17bc
Eucalyptus oil (2%)	18.78	19.12	20.49	21.20	22.32	23.48	26.14	25.78ab	25.89a	29.01bcd
CaCl ₂ (1%)	18.40	19.39	21.97	22.90	24.33	24.88	26.42	27.85ab	29.28abc	33.44ab
Bavistin (0.1%)	18.07	18.80	20.04	21.92	23.24	25.93	28.68	28.99ab	30.30ab	34.00ab
Wax (10%)	17.17	18.91	19.51	19.80	21.00	21.79	24.90	24.45b	24.40bc	25.90cd
Wax (10%) + Bavistin (0.1%)	17.68	18.31	19.33	19.89	20.55	21.56	22.69	23.35b	23.47c	24.73d
SEm (\pm)	1.098	1.303	1.360	1.068	1.476	1.484	2.67	1.901	1.939	1.673
LSD (=0.05)	ns	ns	ns	ns	ns	Ns	ns	5.590	5.702	4.921
CV, %	12.3	13.8	13.4	10.1	13.3	12.5	20.7	14.4	13.9	10.9
P-value	0.946	0.987	0.838	0.279	0.450	0.241	0.598	0.073	<0.001	<0.001
Grand mean	17.89	18.87	20.24	21.09	22.25	23.76	25.75	26.35	27.85	30.76

LSD = Least Significant Difference, SEM = Standard Error of Mean, and CV = Coefficient of variation, ns = non-significant

pH of the fruit

None of the treatment had any significant effects on pH of fruits as shown in table 6. Fruits that were untreated showed maximum (3.73 to 4.48) pH followed by Calcium chloride (1%) (4.05 to 4.43) which were statistically at par with Bavistin (0.1%) treated fruits (4.05 to 4.43) up to 41 days of storage. When the storage period was increased, the pH value of Mandarin fruits was also increased gradually under all the treatments. It

may be due to the conversion and utilization of different acids in the respiration process (Rokaya, 2017). While the minimum pH value was retained by the fruits those were coated with paraffin wax 10% in combination with Bavistin 0.1% which might be due to the slower process of respiration and utilization of organic acids present in mandarin fruits which were statistically at par with cinnamon oil 2% treated fruits.

Treatments	pH of fruit juice on days indicated									
	Day 3	Day 6	Day 9	Day 12	Day 15	Day 18	Day 21	Day 24	Day 27	Day 41
Control	3.73b	3.88	3.95	4.13	4.15	4.23	4.35	4.38	4.43	4.48
Cinnamon Oil (2%)	3.98a	4.08	4.10	4.10	4.10	4.18	4.18	4.28	4.38	4.10
Eucalyptus oil (2%)	3.93ab	3.98	4.20	4.20	4.25	4.25	4.30	4.35	4.38	4.38
CaCl ₂ (1%)	4.05a	4.10	4.13	4.15	4.20	4.23	4.25	4.35	4.40	4.43
Bavistin (0.1%)	4.05a	4.08	4.13	4.20	4.23	4.23	4.28	4.30	4.40	4.43
Wax (10%)	4.13a	4.13	4.15	4.23	4.25	4.25	4.25	4.28	4.28	4.30
Wax (10%) + Bavistin (0.1%)	3.90ab	4.00	4.10	4.10	4.13	4.13	4.13	4.13	4.18	4.18
SEm (±)	0.0748	0.0815	0.1173	0.1795	0.1356	0.997	0.959	0.940	0.899	0.1654
LSD (=0.05)	0.2200	ns								
CV, %	3.8	4.0	5.7	8.6	6.5	7.2	7.3	7.4	6.9	7.6
P-value	0.025	0.368	0.847	0.997	0.972	0.997	0.959	0.940	0.899	0.635
Grand mean	3.96	4.03	4.11	4.16	4.19	4.21	4.25	4.29	4.35	4.33

LSD = Least Significant Difference, SEM = Standard Error of Mean, and CV = Coefficient of variation, ns = non-significant

Postharvest life

Fruits treated with Wax (10%) and Bavistin (0.1%) showed maximum postharvest life of 73 days followed

by Wax (10%). Similarly, minimum span was recorded in control fruits (46days) followed by Eucalyptus oil (2%) (51 days) which is shown in the diagram below.




Figure 1: The Postharvest life of different treatments under ambient room condition

Conclusion

Prolongation of shelf life, as well as the quality of mandarin fruit, could be retained with the use of different surface coatings than without using them. Paraffin wax coating prolonged the storability of mandarin fruits up to 73 days without adversely affecting their physico-chemical and chemical parameters. Hence, coating of mandarin fruits with paraffin wax may be useful for extending their shelf life and effective in stabilizing the market demand. (Even if we used Bavistin in the surface of the fruit and at low concentration as positive control (for experimental purpose), sometimes its use may be harmful if used haphazardly, so we should avoid using such chemicals in real life scenario).

References

- Ahmad, S., Thakur, K., & Siddiqui, M. W. (2013). Postharvest treatments for preserving quality of 'kinnow' fruit under different storage conditions. *Advances in Horticultural Science*, 1-6.
- Akhtar, A., Abbasi, N. A., & Hussain, A. (2010). Effect of calcium chloride treatments on quality characteristics of loquat fruit during storage. *Pakistani Journal of Botany*, 181-188.
- Bhattarai, R. R., Rijal, R. K., & Mishra, P. (2013). Post-harvest losses in mandarin orange: A case study of Dhankuta District Nepal. *African Journal of Agricultural Research*, 763-767.
- Bisen, A., Pandey, S., & Patel, N. (2012). Effect of skin coatings on prolonging shelflife of kagzi lime fruits (*Citrus aurantifolia* swingle). *Journal of Food Science and Technology*, 753-759.
- FAO. (2016, June 3). Statistical Year Book. Retrieved from Food and Agriculture Report: <http://www.faostat.fao.org/site/339/default.aspx>
- Hassan, Z., Lesmayati, S., Qomariah, R., & Hasbianto, A. (2014). Effects of wax coating application and storage temperatures on the on the quality of tangerine citrus (*Citrus reticulata*) var. Siam Banjar. *International Food Research Journal*, 641-648.
- Jholgiker, P., & Reddy, B. (2007). Effect of different surface coating material on postharvest physiology of (*Annona squamosa* L.) fruits under ambient and zero energy cool chamber storage. *Indian Journal of Horticulture*, 64(1), 41-44.
- Kader, A. (2005). Increasing food availability and reducing postharvest losses of fresh produce. In *Proceeding of 5th International postharvest symposium*. F. Mencarelli & P. Tonutti, Acta Horticulture.
- Mahajan, B., Dhatt, A., & Sandhu, K. (2005). Effect of different postharvest treatments on the storage life of kinnow. *Journal of Food Science and Technology*, (Mysore) 42.4: , 296-299.
- Mahajan, B., Dhillon, W., & Kumar, M. (2013). Effect of surface coatings on the shelflife and quality of kinnow fruits during storage. *Journal of Postharvest Technology*, 008-015.
- MoAD. (2016). *Statistical Information and Nepalese Agriculture*. Singhdurbar, Kathmandu, Nepal: Agri Statistics Section, Agribusiness Promotion and Statistics Division, Ministry of Agriculture Development.
- Parashar, M. (2009). Post'-harvest profile of mandarin. Government of India, Ministry of Agriculture, (Department of Agriculture & Cooperation), Directorate of Marketing & Inspection, Branch Head Office Nagpur. Nagpur, India. pp100.
- Rokaya, P. R. (2017, December). Effect of altitude and various pre and postharvest factors on quality and shelflife of mandarin (*Citrus reticulata*, Blanco) (Ph.D. thesis). Agriculture And Forestry University, Rampur, Chitwan, Nepal.
- Sahu, B. (2016, July). Effect of different postharvest treatments on prolonging shelflife of sugar apple (*Annona squamosa* L.) (M.Sc. Thesis). Indira Gandhi Krishi Vishwavidyalaya, India.
- Shahid, M., & Abbasi, N. (2011). Effect of bee wax coatings on physiological changes in fruits of sweet orange cv. "Blood Red". *Sarhad Journal of Agriculture*. 27(3), 385-394.
- Statistical Year Book. (2015/16). Government of Nepal, National Planning Commission, Central Bureau of Statistics, Thapathali, Kathmandu, Nepal.
- Waskar, D., & Gaikwad, R. (2005). Effect of various postharvest treatments on extension of shelflife of keshar mango fruits. *Indian Journal of Agriculture Research*, 95-102.