

# Clinical profile of PCOS patients at a tertiary care center in Gandaki Province of Nepal



Mahendra Raj Pandey

Lecturer, Department of Obstetrics and Gynaecology, Manipal College of Medical Sciences, Pokhara, Nepal

Submission: 06-06-2020

Revision: 07-07-2020

Publication: 01-09-2020

## ABSTRACT

**Background:** Polycystic Ovary Syndrome is one of the most common cause of anovulation in women of reproductive age, with a prevalence of upto 10%. Oligomenorrhea, hirsutism and obesity together with enlarged polycystic ovaries are the diagnostic criteria of PCOS. These women are vulnerable to type II diabetes, dyslipidemia, premature arteriosclerosis, and endometrial carcinoma. **Aims and Objectives:** The aim and objective of this study was to see the various clinical profile of PCOS patients at a Medical College in Nepal and to make patients aware about the long term sequelae associated with it. **Materials and Methods:** This was a retrospective study conducted in the outpatient department of Obstetrics and Gynaecology at Manipal Teaching Hospital from January 2018 till June 2019. A total of 90 patients presenting with oligomenorrhoea, obesity, acne, infertility and hirsutism were included in the study. The exclusion criteria were young women who had their menarche less than 2 years back, women older than 45 years, and patients on exogenous estrogen or progesterone therapy. **Results:** Out of 90 patients studied, 37(41%) were married and 53(59%) were unmarried. There were many patients from urban compared to rural areas (67% vs 33%). Many patients were either overweight (22%) or obese (61%) and 6% of the patients were underweight. Mean age at the time of presentation was 31.66 years with many patients (37%) between 30-35years age group. Majority of the patients were either oligomenorrhoeic or amenorrhoeic (78 %). Weight gain was present in 83% of the patients. Infertility was present in 81% of the married patients. Fifteen women (17%) had hypothyroidism and were receiving treatment for the same. Ultrasonography was done in all the patients and typical necklace pattern of follicular arrangement was seen in 70 (78%) patients. **Conclusion:** The diagnosis of PCOS can be reliably predicted with the help of clinical features and sonography in women according to Rotterdam criteria. Oligomenorrhoea or amenorrhoea and weight gain were main symptoms in our patients. It is important to educate and make unmarried PCOS patients aware of its association with infertility and long-term medical problems.

**Key words:** PCOS; Anovulation; Oligomenorrhea; Obesity

## INTRODUCTION

Polycystic Ovary Syndrome is one of the most common cause of anovulation in women of reproductive age, with a prevalence of upto 10%.<sup>1</sup> PCOS is a growing problem among Nepalese women with a wide variety of presentations. In 1935 Irving F. Stein and Michael L. Leventhal described a symptom complex due to anovulation.<sup>2</sup> Oligomenorrhea, hirsutism and obesity together with enlarged polycystic ovary were the diagnostic criteria of PCOS.

It is a complex clinical presentation and consists of a triad of oligomenorrhea, hirsutism and obesity, and

is now recognized as a heterogeneous disorder that results in overproduction of androgens, primarily from the ovary, and is associated with insulin resistance. The first recognition of an association between glucose intolerance and hyperandrogenism was the famous report of the bearded diabetic woman by Archard and Thiers in 1921.<sup>1</sup>

PCOS may present with amenorrhoea, infertility, features of hyperandrogenemia, signs of metabolic disturbances like insulin resistance, and dyslipidemia. These women are vulnerable to type II diabetes, dyslipidemia, premature arteriosclerosis, and endometrial carcinoma.<sup>3,4</sup>

### Access this article online

**Website:**

<http://nepjol.info/index.php/AJMS>

**DOI:** 10.3126/ajms.v11i5.29325

**E-ISSN:** 2091-0576

**P-ISSN:** 2467-9100

Copyright (c) 2020 Asian Journal of Medical Sciences



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

### Address for Correspondence:

Dr. Mahendra Raj Pandey, Lecturer, Department of Obstetrics and Gynaecology, Manipal College of Medical Sciences, Pokhara, Nepal.

**Mobile:** +977-985603092, **E-mail:** Sarkar\_mrp@yahoo.com

## MATERIALS AND METHODS

This was a cross-sectional study conducted in the outpatient department of Obstetrics and Gynaecology at Manipal Teaching Hospital from January 2018 till June 2019 over 18 months period. A total of 90 patients presenting with oligomenorrhoea, obesity, acne, infertility and hirsutism were included in the study. The exclusion criteria were young women who had their menarche less than 2 years back, women older than 45 years, and patients on exogenous estrogen or progesterone therapy. Presence of at least two criteria from clinical, hormonal, and abdominal sonography were considered diagnostic of PCOS. Women with complaints of oligomenorrhoea or amenorrhoea, signs or symptoms of hyperandrogenism, abdominal ultrasonography showing at least 12 follicles (2-9 mm in diameter) arranged peripherally around a dense core of ovarian stroma or scattered throughout an increased amount of stroma were enrolled in the study. Detailed menstrual history, marital status, and parity were recorded. For the purpose of this study, all diagnoses were made by a consultant gynecologist in association with a senior radiologist. The study was approved by Institutional Review Committee.

Classification of BMI: <18.5 kg/m<sup>2</sup>: underweight, normal BMI: 18.5-24.9 kg/m<sup>2</sup>, Overweight: BMI 25.0-29.9 kg/m<sup>2</sup>, class I obesity: BMI 30-34.9 kg/m<sup>2</sup>, class II obesity: BMI 35-39.9 kg/m<sup>2</sup> and class III obesity was considered as BMI ≥ 40kg/m<sup>2</sup>.<sup>5</sup>

Hirsutism was scored according to modified Ferriman Gallwey score.<sup>6</sup> Grading of severity was based on the score assessed as <4 - mild, 4-7 as moderate and ≥8 - severe.

The data was entered and analysis was performed using statistical package for the social sciences (SPSS, version 25). Mean value and percentages were calculated for various factors.

## RESULTS

Out of 90 patients studied, 37(41%) were married and 53(59%) were unmarried. There were many patients from urban compared to rural areas (67% vs 33%).

Many patients were either overweight (22%) or obese (61%) and 6% of the patients were under weight.

Mean age at the time of presentation was 31.66 years with many patients (37%) between 30-35years age group.

Distribution of patients according to symptomatology: Majority of the patients were either oligomenorrhoeic or amenorrhoeic (78 %). Weight gain was present in 83% of the patients and many patients attributed it due to menstrual

**Table 1: Classification of patients according to BMI**

BMI	Numbers	%
<18.5; underweight	5	6
18.5-24.9; Normal	10	11
25-29.9; Overweight	20	22
>30: Obese	55	61

**Table 2: Age distribution of patients**

Age group	Numbers	Percentage
20-25	10	11
25-30	23	26
30-35	33	37
35-40	20	22
40-45	4	4

**Table 3: Distribution of patients according to symptoms**

Symptoms	Numbers	Percentage (%)
Oligo or Amenorrhoea	70	78
Infertility (out of 37)	30	81
Hirsutism	20	22
Acne	25	28
Acanthosis Nigricans	30	33
Weight gain	75	83

problems. Infertility was present in 81% of the married patients. Hirsutism was present in 22% patients and acne was present in 28% patients. Acanthosis Nigricans was present in 33% patients which is a feature of insulin resistance.

Fifteen women (17%) had hypothyroidism and were receiving treatment for the same. Seven women (8%) had type 2 diabetes mellitus and were on oral hypoglycemics and five patients were on treatment for hyperprolactinaemia. Four patients had co-existing hypertension.

Ultrasonography was done in all the patients and typical necklace pattern of follicular arrangement was seen in 70 (78%) patients.

## DISCUSSION

In this study, we have assessed the clinical presentation among the women diagnosed with PCOS according to the Rotterdam criteria in Gandaki province of Nepal. In our study, majority of the patients were less than 35 years indicating that PCOS is becoming more prevalent in younger population. It was more prevalent among urban people which could be attributed to lack of awareness among rural people and lack of gynaecologists and radiologists in rural areas to diagnose such patients. Normal ovulation results in regular menstrual cycles (28-35 days cycle). Anovulation is the pathognomic feature of PCOS and

results in irregular menstrual cycles. Therefore, persistent menstrual irregularities (resulting from anovulation) seem to be better predictor compared to biochemical parameters. Oligomenorrhea is one of the diagnostic criteria of PCOS. In the present study 78% patients complained of oligomenorrhea or amenorrhoea [Table 3]. Balen et al have reported that menstrual disturbances were seen in 72% of the PCOS women which is comparable to our study. Among them oligomenorrhoea (60%) was the commonest followed by amenorrhoea (11%).<sup>7</sup> In the present study, 81% married PCOS women complained of infertility [Table 3]. Pfeifer SM have also mentioned infertility as one of the long-term sequelae of PCOS.<sup>8</sup> Obesity is common in women who have PCOS. It increases the risk of metabolic dysfunction. Insulin resistance is worsened by presence of obesity. The prevalence of obesity in these women varies. Obesity is not essential to make the diagnosis of PCOS.<sup>9</sup> The results of present study show that PCOS was present in both obese (83%) and non-obese (17%) subjects. In developing nations, obesity is mainly caused by sedentary lifestyle, physical inactivity and consumption of high calorie rich foods. Nepal is currently observing a rise in the number of individuals in the middle-class who are detected with obesity. Obesity is highly observed in the PCOS patients.<sup>10</sup> Lim et al in a systemic review and meta-analysis concluded that women with PCOS had a greater risk of overweight, obesity and central obesity.<sup>11</sup> The drawback of this study was that it was done in a small sample size and hormonal analysis could not be done in all the subjects and it was not included in the study and clinical and radiological parameters were considered for the diagnosis of PCOS.

## CONCLUSION

The diagnosis of PCOS can be reliably predicted with the help of clinical features and sonography in women according to Rotterdam criteria. Oligomenorrhoea or amenorrhoea and weight gain were main symptoms in our patients. Prevalence of PCOS in infertile population was significantly high. Though obesity is common in PCOS, non-obese women are also at risk of PCOS. There is a need to increase awareness regarding role of obesity as well as of acanthosis nigricans as very few patients were aware of their abnormal BMI and skin discoloration. PCOS is becoming a major cause of infertility in women. It is important to

educate and make unmarried PCOS patients aware of its association with infertility and long-term medical problems.

## ACKNOWLEDGMENTS

The author would like to thank all patients inducted during the entire study duration.

## REFERENCES

- Dunaif A. Insulin resistance and the polycystic ovary syndrome: Mechanism and implications for pathogenesis. *Endocr Rev.* 1997; 18: 774-800. <https://doi.org/10.1210/er.18.6.774>
- Stein IF and Leventhal ML. Amenorrhea associated with bilateral polycystic ovaries. *Am Journal of Obstet and Gynecol.* 1935; 29:181. [https://doi.org/10.1016/S0002-9378\(15\)30642-6](https://doi.org/10.1016/S0002-9378(15)30642-6)
- Hardiman P, Pillay OC and Atiomo W. Polycystic ovary syndrome and endometrial carcinoma. *Lancet.* 2003; 361(9371): 1810-1812. [https://doi.org/10.1016/S0140-6736\(03\)13409-5](https://doi.org/10.1016/S0140-6736(03)13409-5)
- Balen A. Polycystic ovary syndrome and cancer. *Human Reprod. Update* 2001; 7:522-525. <https://doi.org/10.1093/humupd/7.6.522>
- BMI classification. World Health Organization. Retrieved 15 February 2014.
- Hatch R, Rosenfield RL, Kim MH and Tredway D. Hirsutism: Implications, etiology, and management. *Am Journal of Obstet and Gynecol.* 1981; 140:815-830. [https://doi.org/10.1016/0002-9378\(81\)90746-8](https://doi.org/10.1016/0002-9378(81)90746-8)
- Balen AH, Conway GS, Kaltas G, Techatrasak K, Manning PJ, West C, et al. Polycystic ovarian syndrome: The spectrum of disorder in 1741 patients. *Hum Reprod.* 1995; 10:2107-2111. <https://doi.org/10.1093/oxfordjournals.humrep.a136243>
- Pfeifer SM and Kives S. Polycystic ovary syndrome in the adolescent. *Obstet Gynecol Clin North Am.* 2009; 36:129-135. <https://doi.org/10.1016/j.ogc.2008.12.004>
- Kalra A, Nair S and Rai L. Association of obesity and insulin resistance with dyslipidemia in Indian women with polycystic ovarian syndrome. *Indian J Med Sci.* 2006; 60:447-453. <https://doi.org/10.4103/0019-5359.27971>
- Ehrmann D. Medical progress: Polycystic ovarian syndrome. *The New England Journal of Medicine.* 2005; 352:1223-1236. <https://doi.org/10.1056/NEJMr041536>
- Lim SS, Davis MJ, Norman RJ and Moran LJ. Overweight, obesity and central obesity in women with polycystic ovarian syndrome: A systemic review and meta-analysis. *Hum Reprod update* 2012; 18 (6):618-637. <https://doi.org/10.1093/humupd/dms030>

### Authors Contribution:

MRP- Concept and design of the study, literature review, data collection and its analysis, manuscript writing and revision of manuscript.

### Work attributed to:

Department of Obstetrics and Gynaecology, Manipal College of Medical Sciences and Hospital, Phulbari campus, Pokhara, Nepal.

### Orcid ID:

Dr. Mahendra Raj Pandey- <https://orcid.org/0000-0002-0066-7652>

**Source of support:** None. **Conflicts of Interest:** The author declares there is no conflict of interest.