

# COVID-19 Vaccine Acceptance and Its Associated Factors Among Young Adults

Shruti Regmi,<sup>1</sup> Sanchita Subedi,<sup>1</sup> Parbati Datheputhe,<sup>1</sup> Sita Rai Koirala<sup>1</sup>

<sup>1</sup>Janamaitri Foundation Institute of Health Sciences, Hattiban, Lalitpur, Nepal.

## ABSTRACT

### Background

COVID-19 pandemic, which has rapidly spread throughout the globe in several waves and mutations, was initially identified in Wuhan, China, in late 2019. Vaccines against this deadly virus were developed quickly in about a year. Despite the availability of vaccines, acceptance of the vaccines among the general population is influenced by various factors and are important to be identified for a successful vaccination program. The objective of this study was to evaluate the COVID-19 vaccine acceptance among young adults and its associated factors.

### Methods

Nepalese young adults were surveyed online using a descriptive cross-sectional study approach. A bilingual, self-administered questionnaire consisting of six parts-socio-demographic information, health-related information, knowledge and attitude/ beliefs about COVID-19, perceived threat to COVID-19, beliefs about COVID-19 vaccination, and vaccine acceptability was used to collect data. The collected data was analysed using SPSS version 20.

### Results

Out of the 467 participants surveyed, most of them (94.2%) were willing to take the vaccine. Among them, 80.5% were willing to take it immediately. The main reason for COVID-19 vaccine refusal was because of doubt about the safety of the COVID-19 vaccine. The regression results showed that healthcare-related professionals/students and perceived benefits of the COVID-19 vaccine were significantly associated with vaccine acceptance.

### Conclusions

This study reflected the high acceptance of COVID-19 vaccination among Nepalese adults during the pandemic. Concerned authorities and health professionals should emphasize addressing concerns about vaccine safety and disseminating information about the benefits of vaccines to overcome vaccine hesitancy.

**Keywords:** attitude/beliefs; COVID-19 vaccine; knowledge; perceived threat; vaccine acceptance.

## INTRODUCTION

COVID-19 was first identified in Wuhan, China, and has since rapidly spread worldwide as a pandemic in various waves and variants of mutation.<sup>1,2</sup> Nepal reported its first verified case on 23<sup>rd</sup> January 2020, with more than 612,000 confirmed cases and 8506 deaths by 15<sup>th</sup> June, 2021.<sup>3,4</sup> This pandemic has a devastating effect on education, agriculture, politics, and economic growth in addition to physical and mental health.<sup>5,6</sup> COVID-19 vaccines were developed

in less than a year as the safest way to control this pandemic.<sup>7</sup> However, vaccine acceptance is context-specific, directly affecting the success of the vaccination program.<sup>8,9</sup> The systemic review showed a wide range of variation across the countries, with the highest COVID-19 vaccine acceptance rate in Ecuador (97.0%) and the lowest in Kuwait (23.6%).<sup>10</sup> The issues of rapid development of COVID-19 vaccines, identification of new variants of coronavirus associated with increased transmissibility

**Correspondence:** Shruti Regmi, Janamaitri Foundation Institute of Health Sciences, Hattiban, Lalitpur. Email: shrutiregmi27@gmail.com, Phone: +977-9815255772. **Article received:**2023-05-21. **Article accepted:**2024-01-05.

and death, re-infection with COVID-19, and post-vaccine infections seem to bear causing significant concerns.<sup>7,11-13</sup> Hence, this study aims to assess vaccine acceptance by Nepalese adults with its associated factors and explore the reasons for vaccine refusal.

## METHODS

An analytical cross-sectional study design was used to conduct an online survey among Nepalese adults of aged 18 years or above by using social media platform (Facebook, Viber, Twitter, What's App) or email, had not received the COVID-19 vaccine, and were willing to participate in the study with informed consent were selected as study participants. Ethical permission was obtained from Nepal Health Research Council (NHRC) (Reg. No. 391/2021) to conduct this study. Participants were automatically directed to the informed consent page followed by the survey questionnaire, which was presented in bilingual form (English and Nepali), after clicking the survey link. The sample size (n) was determined using the Cochran formula,<sup>14</sup> assuming that 50% of general adults have a factor of interest in COVID-19 vaccines and 5% allowable error at a 95% confidence interval (CI). Sample size was 384 and after taking 20% non-response rate, **sample size was 461 but this study was conducted 467**. Participants were selected using the snowball sampling technique. The data was collected using a self-administered survey questionnaire developed in Google Forms based on previous studies to assess vaccine acceptance for COVID-19 and its associated factors.<sup>15-18</sup> The research instrument consisted of six parts: part I consist of socio-demographic information, part II consist of Health-related information (consisted of six questions), part III: Knowledge and attitude/ beliefs about COVID-19, part IV consist of perceived threat to COVID-19, part V: Beliefs about COVID-19 vaccination. Responses to part IV and part V were categorized as (3) Agree, (2) Neutral, and (1) Disagree. Negatively phrased items were recoded. Part VI: Questions related to vaccine acceptability. The instrument was then pre-tested on 46 participants (10% of the sample size) who were contacted through email. For clarity and

convenience, both English and Nepali versions of the questionnaire were made available for data collection. The final dataset was imported and analyzed using Statistical Package for the Social Sciences (SPSS) version 20. The data were analyzed using descriptive and inferential statistics. Descriptive statistics such as frequency and percentage for categorical variables; and mean and standard deviation for continuous variables were used. Inferential statistics such as the Chi-square test and Fisher's exact test were used to find the association between independent variables with COVID-19 vaccine acceptance considering p-value <0.05 as significant. Then, logistic regression was used for finding predictors (factors) of COVID-19 vaccine acceptance considering a 95% confidence interval.

## RESULTS

A total of 467 individuals took part in this survey. More than two-thirds of the participants (68.3%) were between the ages of 21 and 30 years, with individuals over the age of 40 constituting only 2.8% of the study population. Similarly, more than half of the participants (59.5 %) were female, and the majority (78.8%) were unmarried. Likewise, most of the participants (89.7%) followed the Hindu religion, 77.7% were residing in Bagmati province, and 55% were from the Hilly region. More than one-third of the participants (35.5%) were related to the healthcare profession (Table 1). A large proportion of the participants (74.5%) reported having good health and no major illnesses/ comorbidities (94%). More than half of the participants (i.e., 52.2% and 53.1%) were not infected with COVID-19 and had not been exposed to a confirmed case of COVID-19 respectively. whereas, more than half of the participants (54.8%) responded that their family members were infected with COVID-19. Most of the participants (89.7%) had never received an influenza vaccine (Table 2).

Most of the participants (94.2%) were willing to vaccinate with the COVID-19 vaccine. Among them, 80.5% were willing to take it immediately, and the rest (19.5%) wanted to delay. The majority of them

(65.5%) expressed that they would pay for a vaccine out of pocket if the government did not provide it for

Table 1. Socio-demographic characteristics of the participants. (n=467)	
Socio-demographic characteristics	Frequency (%)
<b>Age (in completed years)</b>	
20	104 (22.3)
21-30	319 (68.3)
31-40	31(6.6)
>40	10 (2.8)
Mean age ± S.D	24.04 ± 5.8
<b>Sex</b>	
Female	278(59.5)
Male	189(40.5)
<b>Religion</b>	
Hinduism	419(89.7)
Buddhism	33(7.1)
Christianity	10(2.1)
Atheism	4(0.9)
Islam	1(0.2)
<b>Currently residing Province</b>	
Province 1	37(7.9)
Province 2	14(3)
Bagmati Province	363(77.7)
Gandaki Province	34(7.3)
Lumbini Province	12(2.6)
Karnali Province	3(0.6)
Sudurpaschim Province	4(0.9)
<b>Ecological Region</b>	
Terai Region	201(43.1)
Hilly Region	257(55)
Mountain Region	9(1.9)
<b>Marital status</b>	
Unmarried	368 (78.8)
Married	98 (21)
Divorced	1(0.2)
<b>Educational Level</b>	
No formal education	2(0.4)
Primary level	2(0.4)
Secondary level	10(2.1)
Higher Secondary level	105(22.5)
University level	348(74.5)
<b>Employment status</b>	
Unemployed	336(71.9)
Employed	131(28.1)
<b>Healthcare professionals/ students</b>	
Yes	166(35.5)
No	301(64.5)

free (Figure 1).

Regarding the reasons given by the 27 participants who refused to accept the COVID-19 vaccine,

Table 2. Health-related information of the participants. (n=467)	
Characteristics	Frequency (%)
<b>Self-reported health status</b>	
Excellent	96(20.6)
Good	348(74.5)
Poor	23(4.9)
<b>Presence of major illness/comorbidity</b>	
No	439(94)
Yes	28(6)
<b>Self-infected with COVID-19</b>	
No	244(52.2)
Yes	74(15.8)
Maybe/not tested	149(31.9)
<b>Exposed with a confirmed case of COVID-19</b>	
No	248(53.1)
Yes	219(46.9)
<b>Family member infected with COVID-19</b>	
No	211(45.2)
Yes	256(54.8)
<b>History of receiving Influenza vaccine</b>	
No	419(89.7)
Yes	48(10.3)
<b>Live with vulnerable/high-risk group</b>	
Yes	229(49)
No	238(51)




Figure 1. Participants' Acceptability for COVID-19 Vaccine.

51.9% were doubtful about its safety, 40.7% were concerned about its efficacy, and nearly 15% believed vaccination was unnecessary because they did not trust them (Figure 2).




Figure 2. Reasons for Vaccine Refusal (n=27).

Table 3. Association between Willingness to get COVID-19 Vaccination with the Different characteristics of Participants. (n= 467)			
Characteristics	Willing to get COVID-19 vaccination		$\chi^2$ (p-value)
	No	Yes	
<b>Age</b>			
Less than 25	17	307	0.56
Equal to or above 25	10	133	-0.46
<b>Sex</b>			
Male	15	174	2.7
Female	12	266	-0.1
<b>Province</b>			
Bagmati Province	21	342	0
Other	6	98	-0.99
<b>Ecological region</b>			
Terai	9	192	1.1
Hilly/Mountain	18	248	-0.29
<b>Marital status</b>			
Unmarried	19	350	1.29
Married	8	90	-0.26
<b>Religion</b>			
Hinduism	27	392	-
Other	0	48	-0.1
<b>Education</b>			
University level	16	332	3.51
Below University level	11	108	-0.06
<b>Employment status</b>			
Employed	14	117	8.04
Unemployed	13	323	(0.005*)
<b>Profession</b>			
Health care	1	165	12.68
Other than health care	26	275	(< 0.001*)
<b>Self-reported health status</b>			
Poor	0	23	-
Good to Excellent	27	417	-0.39
<b>Major illness/Co-morbidities</b>			
Absent	23	416	-
Present	4	24	-0.07
<b>Self-Infected with COVID-19</b>			
No/not tested	22	371	-
Yes	5	69	-0.6
<b>Family member infected with COVID-19</b>			
No	15	196	1.25
Yes	12	244	-0.26
<b>Exposure with confirmed case of COVID-19</b>			
No	10	238	2.97
Yes	17	202	-0.09
<b>History of Influenza vaccination</b>			
No	24	395	-
Yes	3	45	-0.75
<b>Knowledge level</b>			
Poor	4	43	-
Good	23	397	-0.34
<b>Attitude level</b>			
Negative	10	79	6.01
Positive	17	361	(0.014*)
<b>Perceived threat with COVID-19</b>			
Low	3	19	-
High	24	421	-0.13
<b>Perceived benefit of COVID-19 Vaccine</b>			
Low	7	17	-
High	20	423	(< 0.001*)
<b>Perceived barriers to COVID-19 Vaccination</b>			
Low	2	64	-
High	25	376	-0.4

Note. \*: significant < 0.05

Table 3 shows the bivariate associations between willingness to get the COVID-19 vaccine and the different characteristics of Nepalese adults. Unemployed status, health care professionals/students, positive attitude toward COVID-19, and high perceived benefit of COVID-19 vaccination were significantly associated with higher rates of COVID-19 vaccine acceptance. Table 4 shows the predictors associated with vaccine intention. Those who were related to health care professionals or students were 13.87 times more likely to be willing to receive the COVID-19 vaccine than those who were not (aOR: 13.87; 95%CI: 1.8-106.84, p = 0.012). Participants who perceived the vaccine to have high benefits were 6.73 times more likely to be willing to receive it (aOR: 6.73; 95%CI: 12.281-19.848, p = 0.001).

## DISCUSSION

The present study aims to assess the COVID-19 vaccine acceptance among Nepalese adults with its associated factors. Our study found high acceptance of COVID-19 vaccination among Nepalese adults during the pandemic. Most participants (94.2%) were willing to be vaccinated with the COVID-19 vaccine, and immediately (80.5%) if the vaccine was made available. The result is in accordance with the study conducted in low- and middle-income countries which reported that Nepal had the highest COVID-19 vaccine acceptance rate with an average acceptance rate of 80.3% for lower-middle-income countries.<sup>19</sup> The findings are also consistent with those obtained from the studies conducted in the People's Republic of China<sup>20</sup> and India.<sup>21</sup> The Chinese study found that 91.3% of Chinese adults would accept COVID-19 vaccination once it became available, with 52.2% wanting to be vaccinated as soon as possible, and others (47.8%) delaying vaccination until the vaccine's safety was confirmed.<sup>20</sup> Another study conducted in India stated that 86.3% of the population was willing to be vaccinated. However, only 65.8% of the participants responded that they would get vaccinated as soon as possible if the vaccine became available.<sup>21</sup> The vaccine acceptance (94.2%) of this study is much higher as compared to studies conducted in Bangladesh (74.6%),<sup>22</sup> United

<b>Table 4. (Logistic regression analysis) Predictors associated with intent to vaccinate.</b>			
<b>Characteristics</b>	<b>Crude OR (95%CI)</b>	<b>Adjusted OR (95%CI)</b>	<b>p-value</b>
<b>Employment status</b>			
Unemployed	2.973 (1.357-6.511)	2.207 (0.949-5.134)	0.066
Employed	Ref		
<b>Health care professionals/students</b>			
Yes	15.6 (2.097-116.003)	13.868 (1.800-106.835)	0.012*
No	Ref		
<b>Attitude level</b>			
Positive	2.688 (1.186-6.092)	2.415 (0.992-5.878)	0.052
Negative	Ref		
<b>Perceived benefit of COVID-19 Vaccine</b>			
High	8.709 (3.243-23.388)	6.729 (2.281-19.848)	0.001*
Low	Ref		

Note. \*: significant < 0.05

States (69%),<sup>23</sup> Saudi Arabia (64.7%),<sup>24</sup> Greece (57.7%),<sup>25</sup> Kuwait (53.1%),<sup>26</sup> and Russia (41.7%).<sup>16</sup> This shows that the vaccine acceptance rate varies greatly across the globe. The high level of acceptance for COVID-19 vaccination among Nepalese adults reflected the strong demand for the COVID-19 vaccine. The most common reasons for COVID-19 vaccine refusal among the participants in this study were because of doubts about vaccine safety, vaccine efficacy, and lack of trust. Such concerns about vaccine safety and effectiveness appear to be widespread, as reported by studies conducted in the United States,<sup>23</sup> Russia,<sup>16</sup> Kuwait,<sup>26</sup> and China.<sup>20</sup> Unemployment, healthcare-related professionals/students, positive attitude toward COVID-19, and high perceived benefit of COVID-19 vaccination were found to be significantly associated with higher rates of COVID-19 vaccine acceptance in this study. Furthermore, those who were healthcare professionals or students were 13.87 times more likely to be willing to receive the COVID-19 vaccine. It was identified as one of the significant predictors associated with vaccine acceptance in a bivariate logistic regression analysis (aOR: 13.87; 95%CI: 1.8-106.84, p = 0.012). This is consistent with findings from an Indonesian study, which found a higher acceptance rate among Indonesian healthcare workers.<sup>27</sup> Another study conducted in China showed that the acceptance of healthcare workers was significantly higher than that

of non-health-care workers (76.98 Vs 56.19%).<sup>28</sup> This could be because healthcare professionals have a more in-depth understanding of vaccine properties. This study also found that high perceived benefits of the vaccine were associated with COVID-19 vaccine acceptance (aOR: 6.73; 95%CI: 12.28-19.85, p = 0.001) which is consistent with the findings of many other studies.<sup>29,30</sup> The Malaysian study showed that the perceived benefits construct in the Health Belief Model had the highest intention to take the vaccine.<sup>29</sup> Similarly, in the Hong Kong study, perceived vaccine benefits were positively correlated with vaccine acceptance, along with other Health Belief Model constructs such as perceived severity, cues to action, self-reported health outcomes, and trust in the healthcare system or vaccine manufacturers.<sup>30</sup> This study provided baseline data for ongoing monitoring of public acceptance of COVID-19 vaccination in Nepal. This study examined the factors independently associated with potential vaccine uptake, which could help to understand priority groups in the vaccination campaign to design effective pandemic immunization strategies. However, this study has some limitations. The use of an online survey may limit the sample's representativeness in the present study. We only contacted those with internet access and used social media and email through the snowball sampling technique. Therefore, disparities in response rates were observed across the country. The Bagmati province had the highest response rate. Most young

adults (97.2%) with university-level education (74.5%) represented this study, and no senior citizen participated. Non-respondent bias occurs when participants who do not respond to the survey differ in their exposures/outcomes from those who do. Furthermore, because self-reported data can lead to information bias, the study findings may vary from the actual practice. Since the COVID-19 vaccine does not provide lifelong immunity, further studies are needed to investigate the changes in vaccination acceptance and its associated factors during different periods of the pandemic.

## CONCLUSIONS

This study reflected the high acceptance of COVID-19 vaccination among Nepalese adults during the pandemic. Moreover, the majority were willing to pay for a vaccine out of pocket. The safety concern seemed to be the main reason for the vaccine refusal. Concerned authorities and healthcare professionals should focus on addressing doubts regarding vaccine safety and disseminating information about the benefits of vaccines to overcome vaccination hesitancy.

## REFERENCES

1. World Health Organization. Listings of WHO's response to COVID-19 [Internet]. World Health Organization; Jun 2020 [updated 2020 Dec 28; cited 2021 May 2]. <https://www.who.int/news/item/29-06-2020-covidtimeline>
2. Jha HB. Second wave of COVID-19 getting out of control in Nepal [Internet]. Observer Research Foundation (ORF); 2021. <https://www.orfonline.org/expert-speak/second-wave-of-covid-19-getting-out-of-control-in-nepal/>
3. Bastola A, Sah R, Rodriguez-Morales AJ Lal BK, Jha R, Ojha HC, et al. The first 2019 novel coronavirus case in Nepal. *Lancet Infect Dis* [Internet]. 2020; 20(3): 279–280. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7130048/>. doi:10.1016/S1473-3099(20)30067-0
4. Government of Nepal Ministry of Health and Population [Internet]. 2021 [cited 2021 Jun 16]. [https://covid19.mohp.gov.np/covid/englishSituationReport/60c89030a0365\\_SitRep492\\_COVID-19\\_15-06-2021\\_EN.pdf](https://covid19.mohp.gov.np/covid/englishSituationReport/60c89030a0365_SitRep492_COVID-19_15-06-2021_EN.pdf)
5. Gyanwali GP. Socio-cultural impacts of COVID-19 in Nepal. *Molung Educational Frontier* [Internet]. 2020 Dec 31 [cited 2021 May 11];10:13-6. DOI:10.3126/mef.v10i0.34054
6. Poudel K, Subedi P. Impact of COVID-19 pandemic on socioeconomic and mental health aspects in Nepal. *Int J Soc Psychiatry* [Internet]. 2020;66 (8):748-55. <https://journals.sagepub.com/doi/pdf/10.1177/0020764020942247>
7. Stulpin C. COVID-19 vaccines developed at record speed, but experts say trust the data. *Infect. Dis. News* [Internet]. 2020 Dec 14 [cited 2021 May 14]. <https://www.healio.com/news/infectious-disease/20201214/covid19-vaccines-developed-at-record-speed-but-experts-say-trust-the-data>
8. MacDonald NE, SAGE Working Group on Vaccine Hesitancy. Vaccine hesitancy: Definition, scope, and determinants. *Vaccines* [Internet]. 2015 [cited 2021 May 4]; 33(34):4161-64. <https://doi.org/10.1016/j.vaccine.2015.04.036>
9. Dubé E, Laberge C, Guay M, Bramadat P, Roy R, Bettinger J. et al. Vaccine hesitancy: an overview. *Hum Vaccin Immunother* [Internet]. 2013 [cited 2021 May 4];9(8):1763-73. doi: 10.4161/hv.24657
10. Sallam M. COVID-19 vaccine hesitancy worldwide: A concise systematic review of vaccine acceptance rates. *Vaccines (Basel)* [Internet]. 2021 Feb 16 [cited 2021 May 5]; 9(2). <https://doi.org/10.3390/vaccines9020160>
11. Nepal govt. confirms detection of B. 1. 617. 2 variant of coronavirus months after advent of second wave. *The Himalayan Times* [Internet]. 2021 May 19 [cited 2021 May 27]. <https://thehimalayantimes.com/environment/presence-of-third-new-variant-of-coronavirus-confirmed-by-nepal-government>.

12. Keehner J, Horton LE, Pfeffer MA, Longhurst CA, Schooley RT, Currie JS, et al. SARS-CoV-2 infection after vaccination in health care workers in California. *N Engl J Med* [Internet]. 2021 [cited 2021 Jun 1]; 384(18):1774-75. doi:10.1056/NEJMc2101927
13. Cohen JI, Burbelo PD. Reinfection with SARS-CoV-2: implications for vaccines. *Clin Infect Dis* [Internet]. 2020 Dec 18;ciaa1866. doi: 10.1093/cid/ciaa1866, Epub ahead of print. PMID: 33338197; PMCID: PMC7799323.
14. Cochran, WG. *Sampling Techniques*. 3<sup>rd</sup> ed. New York: John Wiley & Sons; 1977. <https://hwbddocuments.env.nm.gov/Los%20Alamos%20National%20Labs/General/14447.pdf>
15. Shmueli L. Predicting intention to receive COVID-19 vaccine among the general population using the health belief model and the theory of planned behavior model. *BMC Public Health* [Internet]. 2021 Apr 6 [cited 2021 May 25]; 21(1):804. <https://doi.org/10.1186/s12889-021-10816-7>
16. Tran VD, Pak TV, Gribkova EI, Galkina GA, Loskutova EE, Dorofeeva VV, et al. Determinants of COVID-19 vaccine acceptance in a high infection-rate country: a cross-sectional study in Russia. *Pharm Pract (Granada)* [Internet]. 2021 [cited 2021 May 27];19(1):2276. doi:10.18549/pharmpract.2021.1.2276
17. Huynh G, Nguyen TV, Nguyen DD, Lam QM, Pham TN, Nguyen HTN, et al. Knowledge about COVID-19, beliefs and vaccination acceptance against COVID-19 among high-risk people in Ho Chi Minh City, Vietnam. *Infect Drug Resist* [Internet]. 2021 [cited 2021 Jun 2]; 14:1773-80. <https://doi.org/10.2147/IDR.S308446>
18. Al-Qerem WA, Jarab AS. COVID-19 vaccination acceptance and its associated factors among a Middle Eastern population. *Front Public Health* [Internet]. 2021 Feb 10 [cited 2021 May 16];9:632914. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7902782/>. doi: 10.3389/fpubh.2021.632914
19. SolísArce JS, Warren SS, Meriggi NF, Scacco A, McMurry N, Voors M, et al. COVID-19 vaccine acceptance and hesitancy in low- and middle-income countries. *Nat. Med.* 2021;27(8):1385-94. <https://doi.org/10.1038/s41591-021-01454-y>
20. Wang J, Jing R, Lai X, Zhang H, Lyu Y, Knoll MD, Fang H. Acceptance of COVID-19 Vaccination during the COVID-19 Pandemic in China. *Vaccines (Basel)*. 2020 Aug 27;8(3):482. doi:10.3390/vaccines8030482, PMID: 32867224; PMCID: PMC7565574.
21. Khan S, Rahman CKF, C V H, Jose B, Tiwari R, Dhama K. COVID-19 Vaccine Acceptance: Beliefs and Barriers Associated with Vaccination Among the General Population in India. *J J. Exp. Biol. Agric. Sci.* 2020;8:S210-S8. doi:10.18006/2020.8(Spl-1-SARS-CoV-2). S210.S218
22. Abedin M, Islam MA, Rahman FN, Reza HM, Hossain MZ, Hossain MA, et al. Willingness to vaccinate against COVID-19 among Bangladeshi adults: Understanding the strategies to optimize vaccination coverage. *PLoS One*. 2021 Apr 27;16(4):e0250495. doi: 10.1371/journal.pone.0250495, PMID: 33905442; PMCID: PMC8078802.
23. Reiter PL, Pennell ML, Katz ML. Acceptability of a COVID-19 vaccine among adults in the United States: How many people would get vaccinated? *Vaccine*. 2020;38(42):6500-7. doi: 10.1016/j.vaccine.2020.08.043
24. Al-Mohaithef M, Padhi BK. Determinants of COVID-19 Vaccine Acceptance in Saudi Arabia: A Web-Based National Survey. *J Multidiscip Healthc.* 2020;13:1657-63. doi: 10.2147/JMDH.S276771
25. Kourlaba G, Kourkouni E, Maistrelis S, Tsopele C-G, Molocha N-M, Triantafyllou C, et al. Willingness of Greek general population to get a COVID-19 vaccine. *Global Health Research and Policy*. 2021;6(1):3. <https://doi.org/10.1016/j.ghrp.2021.01.001>

- org/10.1186/s41256-021-00188-1
26. Alqudeimat Y, Alenezi D, AlHajri B, Alfouzan H, Almokhaizeem Z, Altamimi S, et al. Acceptance of a COVID-19 Vaccine and its Related Determinants among the General Adult Population in Kuwait. *Med Princ Pract*. 2021;10:2052-61. doi:10.1159/000514636
  27. Harapan H, Wagner AL, Yufika A, Winardi W, Anwar S, Gan AK, et al. Acceptance of a COVID-19 Vaccine in Southeast Asia: A Cross-Sectional Study in Indonesia. *Front Public Health*. 2020;8:3814. doi:10.3389/fpubh.2020.00381
  28. Wang MW, Wen W, Wang N, Zhou MY, Wang CY, Ni J, et al. COVID-19 Vaccination Acceptance Among Healthcare Workers and Non-healthcare Workers in China: A Survey. *Front Public Health*. 2021 Aug 2;9:709056. doi: 10.3389/fpubh.2021.709056, PMID: 34409011; PMCID: PMC8364953.
  29. Wong LP, Alias H, Wong P-F, Lee HY, AbuBakar S. The use of the health belief model to assess predictors of intent to receive the COVID-19 vaccine and willingness to pay. *Human vaccines & immunotherapeutics*. 2020;16(9):2204-14 . doi: 10.1080/21645515.2020.1790279
  30. Wong MCS, Wong ELY, Huang J, Cheung AWL, Law K, Chong MKC, et al. Acceptance of the sCOVID-19 vaccine based on the health belief model: A population-based survey in Hong Kong. *Vaccine*. 2021;39(7):1148-56. doi: 10.1016/j.vaccine.2020.12.083, PMCID: PMC7832076

**Citation:** Regmi S, Subedi S, Datheputhe P, Koirala SR. COVID-19 Vaccine Acceptance and Its Associated Factors Among Nepalese Adults. *JCMS Nepal*. 2024; 20(1): 18-25.