

Incidence of Bacteremia and Septicemia in patients attending in tertiary care center, Nepal

Chaudhary R, Karmacharya S, Shrestha S, Dahal R K, Mishra S K, Banjade N R, Kattel H P, Rijal B P, Sherchand J B, Pokhrel B M

Department of Microbiology, Tribhuvan University Teaching Hospital, Kathmandu, Nepal

Correspondence to: Dr Raina Chaudhary

E-mail: raina_chaudhary@yahoo.co.in

Abstract

Introduction: Bacteremia and septicemia is life threatening condition resulting in major cause of mortality and morbidity. The aim of study was to determine the etiology of bacteremia and septicemia with antibiotic sensitivity profile of those organisms.

Methods: A prospective study was carried out among the suspected cases from both inpatient and outpatient of TUTH from October 2009 – March 2010. Blood samples were collected and processed according to standard methodology.

Results: Out of 2259 samples only 237 (10.49 %) showed bacterial growth. The most common isolates among Salmonella group was Salmonella enterica serotype typhi 71 (29.95%) followed by Salmonella enterica serotype Paratyphi A 45 (18.98%). Among non Salmonella group Pseudomonas aeruginosa 34 (14.34%), Klebsiella pneumoniae 22 (9.28%), Acinetobacter spp 15 (6.32%), Citrobacter spp 5 (2.10%), Escherichia coli 3 (1.26%) while Staphylococcus aureus 34 (14.3%) was most common followed by Enterococcus spp 3 (1.26%), Streptococcus spp 2 (0.84%), Coagulase Negative Staphylococcus 2 (0.84%) and Listeria spp 1 (0.42%) among Gram Positive organisms. AntibioGram revealed Cefotaxime, Ceftazidime, Azithromycin and Chloramphenicol for Salmonella group while for non Salmonella Imipenem, Meropenem and Amikacin as most effective antibiotics while Clindamycin, Ciprofloxacin and Ofloxacin for gram positive.

Conclusion: Gram negative bacteria was the predominant organism causing bacteremia and septicemia. Among them salmonella typhi and salmonella paratyphi were the leading aetiology.

Keywords: Bacteremia, Septicemia,

Introduction

Continuous or transient presence of microorganism within the blood stream is Bacteremia. While its dissemination throughout the body with evidence of systemic responses towards microorganism with variable severity is Septicemia. Though the term Bacteremia and Septicemia are interchangeably being used. Whatever may be the definition of Bacteremia and Septicemia both are always threat to every organ of the body and even affecting the

function of the implanted foreign bodies like heart valve, joints, shunts etc¹.

Bloodstream infection are an important cause of serious morbidity and leading cause of mortality and among the most common healthcare associated infection^{2,3}. Study in United States shows it as 10th leading cause of death⁴. They are associated with the syndrome requiring admission to intensive care unit such as sepsis and septic shock⁵⁻⁷. They are not only considered as leading cause of death also

causing disability, worsening of quality of life especially for millions of people in developing countries⁸⁻¹⁰. Study carried out in United States shows mortality rates as 14 days mortality averaged 26%, 28 days mortality 42%¹¹. It has impact on economy of nation as well again study in US, shows during management of Bacteraemia & Septicaemia they found expense of cost around US\$16 billion per year¹¹.

Culturing the blood sample to reveal the presence of microorganism is a highly specific indicator of bloodstream infection and the results of antimicrobial susceptibility testing may assist in the choice of appropriate antimicrobial therapy for such patient. Furthermore early and rapid administration of antimicrobial therapy to such patient shown to reduce mortality and morbidity^{12,13}. Nowadays bacterial drug resistance is an important problem and due to wide variation in bacteria drug resistance study and report in one region or in a period of time necessarily not true for other region or period of time¹⁴. The resistance pattern are related with series of social, environmental and technological changes¹⁵.

The surveillance of bloodstream pathogen in a hospital is important in monitoring the spectrum of microorganism that invade the bloodstream and the type of organism associated with a particular clinical discipline. Such data is often used to determine empiric antibiotic therapy and also to alert clinician the emerging pathogen that may be threat to the community.

Methods

Prospective type of study was done in Bacteriology Laboratory of TUTH, from October 2009-March 2010. All the suspected cases of bacteremia and septicemia from both outpatient and inpatient were included. Of total 2259 blood and bone marrow samples were processed. During processing 5ml of blood was mixed with 45ml brain heart infusion broth for adult and 1ml blood with 9ml infusion for children. After 24 hours blind subculture was done then subsequent subculture was done after viewing turbidity. The broths were kept till 7 days then discarded after blind subculture.

Antibiotic susceptibility testing¹⁶

Antibiotic susceptibility testing was done using Muller Hinton Agar (MHA) by standard disk diffusion technique of Kirby-Bauer method as recommended by CLSI (Clinical Laboratory Standards Institute). Antibiotic tested were Amoxicillin, Azithromycin, Amikacin, Ampicillin+Sulbactam, Amoxicillin+Clavulanic Acid, Co-timoxazole, Chloramphenical, Ciprofloxacin, Cefotaxime, Ceftazidime, Clindamycin Erythromycin Gentamycin, Imipenem, Meropenem, Ofloxacin Piperacillin, Vancomycin.

Results

A total of 2259 specimens from blood and bone marrow were received in the Microbiology Laboratory for culture and sensitivity from October 2009 to March 2010.




Fig. 1: Pattern of growth in blood and bone marrow (n=2259)

Fig.1 shows that only 10% of blood and bone marrow specimens showed significant bacterial growth.

Table 1 shows that *Salmonella enterica* serotype Typhi (36.41%) was the most predominant organism followed by *Salmonella enterica* serotype Paratyphi A (23.07%), *Pseudomonas aeruginosa* (17.43%), *Staphylococcus aureus* (17.43%).

Table 1: Distribution of bacterial isolates (n=237)

Name of organisms	Number	(Percent)
Salmonella enterica serotype Typhi	71	(29.95%)
Salmonella enterica serotype Paratyphi A	45	(18.98%)
Pseudomonas aeruginosa	34	(14.34%)
Staphylococcus aureus	34	(14.34%)
Klebsiella pneumoniae	22	(9.28%)
Acinetobacter spp	15	(6.3%)
Citrobacter freundii	5	(2.1%)
Enterococcus spp	3	(1.26%)
Escherichia coli	3	(1.26%)
Streptococcus spp	2	(0.84%)
Coagulase Negative Staphylococcus spp	2	(0.84%)
Listeria spp	1	(0.42%)

Table 2 shows distribution of Gram negative and Gram positive bacterial isolates. Gram negative (82%) bacteria were significantly predominant than Gram positive (18%) with P value >0.01%.

Table 2: Comparison of Gram negative and Gram positive bacterial isolates

Gram negative bacterial isolates	N (%)	Gram positive bacterial isolates	N (%)
Salmonella enterica serotype Typhi	71 (29.95%)	Staphylococcus aureus	34(14.34%)
Salmonella enterica serotype Paratyphi A	45 (18.98%)	Enterococcus spp	03(1.26%)
Pseudomonas aeruginosa	34 (14.34%)	Streptococcus spp	02(0.84%)
Klebsiella pneumoniae	22 (9.28%)	Coagulase negative Staphylococcus	02(0.84%)
Acinetobacter spp	15 (6.3%)	Listeria spp	01 (0.42%)
Citrobacter freundii	05 (2.1%)		
Escherichia coli	03 (1.26%)		
Total number of Gram negative isolates	195 (82%)	Total number of Gram positive isolates	42 (18%)

Table 3 shows sensitivity pattern of Salmonella enterica serotype Typhi, majority of Salmonella typhi showed susceptibility towards Azithromycin(100%), Cefotaxime (100%) and Ofloxacin (100%) followed by Ciprofloxacin (97%) and Chloramphenicol (92%). Amoxicillin (83%) found out to be least sensitive.

Table 3: Sensitivity pattern of Salmonella enterica serotype Typhi (n=71)

Antibiotics	Sensitive %
Amoxicillin	83%
Azithromycin	100%
Co-trimoxazole	90%
Chloramphenicol	92%
Cefotaxime	100%
Ciprofloxacin	97%
Nalidixic Acid	21%
Ofloxacin	100%

Table 4 shows the sensitivity pattern of Salmonella enterica Paratyphi A, the majority of Salmonella typhi isolates showed susceptibility towards Azithromycin (100%), Cefotaxime (100%) and Ofloxacin (100%) followed by Chloramphenicol (98%) and Co trimoxazole (98%). Amoxicillin (74%) found out to be least sensitive among the tested antibiotic.

Table 4: Sensitivity pattern of Salmonella enterica serotype Paratyphi A (n=45)

Antibiotics	Sensitive %
Amoxicillin	74%
Azithromycin	100%
Co-tromoxazole	98%
Chloramphenicol	98%
Cefotaxime	100%
Ciprofloxacin	89%
Nalidixic Acid	13%
Ofloxacin	100%

Table 5 shows the sensitivity pattern of the *Pseudomonas aeruginosa*, majority of the isolates showed susceptibility towards Imipenem (97%), Meropenem (97%), Amikacin (94%) and Piperacillin(94%) followed by Chloramphenical (91%) and Cefipime (80%). Ceftazidime (76%) and Ciprofloxacin (59%) found out to be least sensitive.

Table 5: Sensitivity pattern of *Pseudomonas aeruginosa* (n=34)

Antibiotics	Sensitive%
Amikacin	94%
Ceftazidime	76%
Ciprofloxacin	59%
Chloramphenical	91%
Cefipime	80%
Imipenem	97%
Meropenem	97%
Ofloxacin	59%
Piperacillin	94%

Table 6 shows sensitivity pattern of *Klebsiella pneumonia*, majority of the isolates showed sensitivity towards Imipenem (91%), Meropenem(91%) followed by Amikacin (64%). Ceftazidime (18%) and Ciprofloxacin (14%) were found to be least sensitive.

Table 6: Sensitivity pattern of *Klebsiella pneumoniae*(n=22)

Antibiotics	Sensitive %
Amikacin	64%
Co-trimoxazole	14%
Ceftazidime	18%
Ciprofloxacin	14%
Chloramphenical	18%
Imipenem	91%
Meropenem	91%
Ofloxacin	27%

Table 7.shows sensitivity pattern of *Staphylococcus aureus*, majority of the isolates showed sensitivity towards Amikacin (100%), Clindamycin (100%) and Cefotaxime (100%) followed by Ciprofloxacin (97%), Vancomycin (94%) and Co trimoxazole (82%). Amoxicillin (62%) and Amoxicillin +Clavulonic acid (65%) had least sensitivity.

Table 7: Sensitivity pattern of *Staphylococcus aureus* (n=34)

Antibiotics	Sensitive%
Amoxicillin	62%
Amoxicillin+Clavulonic Acid	65%
Amikacin	100%
Co-trimoxazole	82%
Ciprofloxacin	97%
Clindamycin	100%
Cefoxitin	88%
Cefotaxime	100%
Erythromycin	79%
Vancomycin	94%

Table 8. shows sensitivity pattern of *Acinetobacter* spp, majority of the isolates showed sensitivity towards Co trimoxazole (100%), Chloramphenical (100%) followed by Amikacin (93%), Imipenem (93%) and Meropenem (93%). Amoxicillin (73%) showed least activity.

Table 8: Sensitivity pattern of *Acinetobacter* spp (n=15)

Antibiotics	Sensitive %
Amoxicillin	73%
Amikacin	93%
Ampicillin+Sulbactam	80%
Co-trimoxazole	100%
Chloramphenical	100%
Ciprofloxacin	80%
Cefotaxime	87%
Imipenem	93%
Meropenem	93%

Discussions

In this study out of 2259 processed samples only 237 (10%) of showed growth. This result was quite similar with the previous study done by Banjara¹⁹ in the same laboratory in1999 with the positivity of 15 percent. But it was lower than the study result of Kathmandu model hospital by Amatya et al² in 2005 who recovered 23% of growth positive, similarly study done by Karki et al²¹ at Kanti Hospital in 2007- 2008 recovered 4.2% of growth positive.

In this study, Gram negative bacteria constituted the major group of isolates comprising 82 percent. This was similar with the results reported by Mehdinejad et al²² as 86.5% and by Mehta et.al²³ as 80.96%.

According to the present study, *S.enterica* serotype Typhi found out to be major cause of bacteraemia and

septicaemia, which accounted for 30% of all isolates. This had a similarity with the previous study in this Hospital in 1999 by Banjara¹⁹ who reported 38.5 percent.

Present study demonstrated that 100% of isolates were sensitive to Ofloxacin and Azithromycin with the least activity for Amoxicillin of 83 percent.

Present Study showed *Salmonella enterica* serotype Paratyphi A contributed second major cause of bacteraemia & septicaemia in this particular Hospital and accounted 19% of all isolates. This result was higher than previous study done in 1999 by Banjara¹⁹, who reported as 11.4percent. But it was lower than study in Model Hospital by Amatya et al²/ 35.8 percent.

Present study showed *Staphylococcus aureus* as 14.3% of all isolates. This result had similarity with study by Mehta et al²³ who reported 13.86% of isolates of *S. aureus*. But it was lower than previous study by Banjara¹⁹, who reported only 42.1% in 1999 from same Hospital.

In this study, 100% of *Staphylococcus aureus* were sensitivity to Clindamycin and Cefotaxime, 97% of isolates were sensitive to Ciprofloxacin & Ofloxacin with the least sensitivity of 62 percent against Ampicillin.

The present study revealed 9.28% of total isolates as *Klebsiella pneumoniae* This results was in accordance to SENTRY³⁴ surveillance reported 7.3% from US & Canada in 1999. In this study, none of the antibiotics were found to be 100% sensitive among the isolates of *K.pneumoniae*. 91% of *K. pneumoniae* were sensitivity to Imipenem & Meropenem followed by 64% of isolates were sensitive to Amikacin and least sensitivity to Cephlosporine group of antibiotics like Cefotaxime Cefipime & Ceftazidime ranging from 14 to 27 percent.

Acinetobacter spp have been implicated in recent years as important nosocomial pathogen, especially in intensive care setting. Despite their low pathogenic potential they are being reported increasingly as the causal organism of numerous hospitals outbreaks in several countries. In a recent international multicenter study, *Acinetobacter* spp were ranked amongst 10 organisms most commonly causing septicaemia in 18 of 44 large European hospitals³⁶. Therefore, this study was not the exception and showed 6.32% growth for *Acinetobacter* spp

Conclusion

From this study, it becomes clear that Gram negative bacteria were the main causative organisms for the bacteremia and septicemia. Among them *Salmonella Typhi* were the leading cause followed by *Salmonella*

Paratyphi A. Sensitivity pattern is constantly changing even in same set up and even geographical & ethnical variation undoughtably showed change in pattern of sensitivity. Therefore, it emphasized on the need for such type of study at regular interval.

Acknowledgement

We would like to thank all the staff member of Microbiology Department for the generous help during this study.

References

1. Lowell S. Young. Sepsis Syndrome. In: Gerald L. Mandell, John E. Bennett, Raphael Dolin, editors, Principle and Practice of Infectious Diseases, 5th Edition, Philadelphia, Pennsylvania, Churchill Livingstone, 2000; 806-819.
2. Deikma DJ, Beckman SE, Chapin KC, Moral KA, Munson E, Doern GV. Epidemiology and Outcome of nosocomial onset bloodstream infection. *J Clin Microbiol* 2003; 41: 3655-60.
3. Karlowsky JA, Jones ME, Draghi DC, Thomsberry C, Sahn DF and Volturo GAs. Prevalance and antimicrobial susceptibility of bacterial isolates from blood culture of hospitalized patients in the United States in 2002. *Ann Clin Microbial Antimicrobial*; 2004; 10: 3-7.
4. National Center for Health Statistics Health, United States, 1998 with socioeconomic status and Health Chart book. DHHS Publication No (PHS) 98-1232. Hyattsville, Md: NCHS; 1998: 271.
5. Balk R. Severe Sepsis and Septic Shock, Definition, Epidemiology and Clinical manifestations. *Crit Care Clin* 2000; 16: 179-92.
6. Anwer SK, Mustafa S, Pariyani S, Ashraf S and Taufiq KM. Neonatal Sepsis and etiological study. *J Pak Med Assoc* 2000; 50: 91-4.
7. Joshi JJ, Ghole VS and Niphadkar KB. Neonatal Gram Negative Bacteriamia. *Indian J Pediatr* 2000; 67: 27-32.
8. Butta ZA and Yusuf K. Neonatal Sepsis in Karachi: Factor determining outcome and mortality. *J Trop Pediatr* 1997; 43: 65-70.
9. Orrett FA and Sherland SM. Neonatal Sepsis and mortality in a regional hospital in Trinidad: Aetiology and risk factors. *Ann Trop Paediatr* 2001; 21: 20-5.
10. Madsen KM, Schonheydr HC, Kristansen B and Sorensen HT. Secular trends in incidence and

- mortality of bacteremia in Danish Country 1981-1994. *APMIS. Acta Pathol Microbiol Immunol Sci* 1991; 107: 346-52.
11. Bone RC. Towards and Epidemiology and natural history of SIRS (Systemic Inflammatory Response Syndrome). *JAMA* 1992; 268: 3452-5.
 12. Leibovisi L, Konisberge H and Pitlik SD. Bacteremia and Fungemia of Unknown origin in adult. *Clin Infect Dis* 1992; 14: 436-9.
 13. Bhothe RR, Furth Van B and Van Den PJ. Aetiology of Community-Acquired Pneumonia: Prospective study among adult requiring admission to hospital. *Thorax*. 1995; 50: 543-7.
 14. Cohen ML. Epidemiology factors influencing the emergence of antimicrobial resistance. *Ciba Found Symp* 1997; 207: 223-231.
 15. Huang SS, Labus BJ, Samhel MC, Wan DT and Reingald AL. Antibiotic resistance pattern of bacterial isolates from blood in San Francisco country, California 1996-1999. *Emerg Infect Dis* 2002; 8: 195-201.
 16. Clinical and Laboratory Standards Institute. In Performance standards for antimicrobial disk susceptibility testing: seventeenth information supplement. CLSI document M100-S17 (ISBN-56238-625-5); Vol. 27, No.1, 2007.
 17. Franklin C, Liolios L, Pegleg AY. Phenotypic detection of carbapenem-susceptible Metallo-beta-lactamase-producing Gram-negative bacilli in the clinical laboratory. *J Clin Microbiol* 2006; 44: 3139-44.
 18. Hakanen A, Kotilainen P, Jalava J, Sittone A, Huovinem. Detection of decreases fluoroquinolone susceptibility in Salmonellas and validation of nalidixic acid screening test. *J Clin Micro* 1999; 37: 3572-7.
 19. Banjara NR and Pokharel BM. Bacteriology of bacteremia/septicemia at Tribhuvan University Teaching Hospital. *J Nep Assoc Med Lab Sci* 2001; 21:30.
 20. Amatya NM, Shrestha B, Lakhak B. Etiological Agents of Bacteraemia and Antibiotic Susceptibility Pattern in Kathmandu Model Hospital. *J Nep Med Assoc* 2007; 46(167):112-8.
 21. Karki S, Rai GK, Manandhar R. Bacteriology Analysis and Antibiotic Sensitivity Pattern of Blood Culture Isolates in Kanti Children Hospital. *J Nep Paediatr Soc* 2010; 30:94-8.
 22. Mehdinejad M, Khosravi AD and Morvaridi A. Study of Prevalence and Antimicrobial Susceptibility Pattern of Bacteria Isolated from Blood Cultures. *J Bio Sci* 2009; 9:249-53.
 23. Mehta M, Dutta P and Gupta V. AntiMicrobial Susceptibility Pattern of Blood Isolates from a Teaching Hospital in North India. *Jpn J Infect Dis* 2005; 58:174-6.
 24. Watson JP, Pettibone EC. Chloramphenicol and ampicillin resistant Salmonella typhi in Nepal. *J Nep Med Assoc* 1991; 29: 259-61.
 25. Lewis MD, Serichantalergs O, Pitarangsi C, Chuanak N, Manson CJ, Regmi LR, Pandey P, Laskan R, Shrestha CD, Malla S. Typhoid fever: a massive single-point source, Multidrug-resistant outbreak in Nepal. *Clin Infect Dis* 2005; 40:554-61.
 26. Tuladhar NR, Banjada NR, Pokharel BM, Rizal B, Manandhar R, Shrestha S, Shah A, Chaurasia S. Antimicrobial resistant bacterial strains from in-patients of Tribhuvan University Teaching Hospital, Kathmandu. *J-IOM* 2003; 1; 19-26.
 27. Pokharel BM, Koirala J, Dahal RK, Mishra SK, Khadga PK, Tuladhar NR. Multidrug-resistant and extended-spectrum beta-lactamase(ESBL)-producing Salmonella enteric (serotype Typhi and Paratyphi A) from blood isolates in Nepal: Surveillance of resistance and a search for newer alternatives. *Int J infect Dis* 2006; 10: 434-8.
 28. Sharma AK. Antimicrobial Resistance Pattern of Salmonella in Kanti Children's Hospital: Which Drug to Choose? *J Nep Paed Soci* 2006; 26(1): 21-23.
 29. Mishra SK, Sah JP, Shrestha R, Lakhey M. Emergence of Nalidixic Acid Resistant Salmonella : A confounding scene in Antibiotic Armamentum. *J Nep Assoc Med Lab Sci* 2008; 9:61-4.
 30. Khanal B, Sharma SK, Bhattacharya SK, Bhattarai NR, Deb M, and Kanungo R. Antimicrobial Susceptibility Patterns of Salmonella enteric Serotype Typhi in Eastern Nepal. *J Helth Popul Nutr* 2007; 25:82-7.
 31. Enne VI, Livermore DM, Stephens P, Hall LM: Persistence of sulphonamide resistance in Escherichia coli in the UK despite national prescribing restriction. *Lancet* 2001; 357: 1325-1328.

32. Kumari N, Mohapatra TM, Singh YI. Prevalence of Methicillin-Resistant *Staphylococcus aureus* MRSA in a Tertiary-care Hospital in Eastern Nepal. *JNMA*; 47(170): 53-6.
33. Sanjana RK, Shah R, Chaudhary N, Singh Y I. Prevalence and antimicrobial susceptibility pattern of methicillin-resistant *Staphylococcus aureus* (MRSA) in CMS teaching hospital: a preliminary report. *J Coll Med Sci-Nep* 2010; 6:1-6.
34. Gales AC, Jones RN, Pfaller MA, Gordon KA, Sader HS. Two year assessment of pathogen frequency and antimicrobial resistance pattern among organisms isolated from skin and soft tissue infections in Latin America hospitals: results from the SENTRY antimicrobial surveillance programme, 1997-98, SENTRY study group. *Int J Infect Dis* 2000; 4: 75-84.
35. Kivanc S, Hale T, Funda ET and Hande A. Bloodstream Infections Caused by ESBL-Producing *E.coli* and *K. pneumoniae* : Risk Factors for Multidrug-Resistance. *The Braz J Infect Dis* 2009; 13:403-7.
36. Prashanth K. and Badrinath S. Simplified phenotypic tests for identification of *Acinetobacter* spp and their antimicrobial status. *J Med Microbiol* 2000; 49: 773-8.
37. Souli M., F.V. Kontopidole, E. Koratzanis, A. Antoniadou et.al. In Vitro activity of tigecycline against multiple-drug-resistant, including pan-resistant, Gram-negative and Gram-positive clinical isolates from Greek hospitals. *Antimicrob Agents Chemother* 2006; 50: 3166-9.
38. Mishra SK, Acharya J, Katel HP, Shah A, Rijal BP, Pokhrel BM. Metallo-beta-lactamase producing bacterial isolates: First report from Nepal from Nepal. *Research Conference on Health Science*. 2009 March 2; 57.