

Response of Wheat Genotypes to Different Levels of Nitrogen

Shukra Raj Shrestha^{1*}, Sarita Manandhar¹, Bedanand Chaudhary¹, Bibek Sapkota², Rudra Bhattarai¹ and Surya Prasad Adhikari¹

¹Regional Agricultural Research Station, Sunsari

²Socio-economics and Agricultural Research Policy Division, Khumaltar, Nepal

*Correspondence: shukrarajshrestha@gmail.com

Received January 2016; Revised September 2016; Accepted October 2016

Scientific Editors: Yagya G. Khadka, Bhava P. Tripathi

Copyright ©2016 NARC. Permits unrestricted use, distribution and reproduction in any medium provided the original work is properly cited

ABSTRACT

A field experiment was conducted for two years using six genotypes of wheat (*Triticum aestivum* L.) for response to different levels of nitrogen (N) use. The experiment was laid out in split plot design with four levels (0, 50, 100 and 150 kg N ha⁻¹) as main plots and six wheat genotypes (BL 3623, BL 3629, BL 3872, NL 1008, NL 1055 and Vijay, a check variety) as sub-plots. Grain yield and other yield components increased linearly in response to N concentrations in both seasons. Only two parameters: days to heading (DOH) and days to maturity (DTM) varied significantly ($p \leq 0.05$) among wheat genotypes in both the years. None of the parameters showed interaction effects in both seasons. Vijay showed highest grain yield of 3.12 t ha⁻¹ in 2013 with the application of 100 kg N ha⁻¹, and 3.23 t ha⁻¹ in 2014 with 150 kg N ha⁻¹. Spike length, productive tillers m⁻², number of spikes m⁻² and test weight were greater with higher N rates. The straw yield of wheat fertilized with 150 kg N ha⁻¹ was the highest in Vijay (4.35 t ha⁻¹) and BL 3872 (4.33 t ha⁻¹), respectively. Vijay with 100 kg N ha⁻¹ produced the highest number of productive tillers m⁻² (276.33) in 2013 and 296.00 with the application of 150 kg N ha⁻¹ in 2014.

Keywords: Nitrogen levels, wheat genotypes, yield

सारांश

गहुँका ६ वटा उन्मोचन योग्य जातहरूमा माटोमा नाइट्रोजनको विभिन्न मात्रा राखी दुई वर्ष सम्म उत्पादन र उत्पादकत्वको अध्ययन गरियो। उक्त परीक्षण Split Plot Design मा संचालन गर्दा मुख्य प्लटहरूमा नाइट्रोजनको चारवटा मात्राहरू (०, ५०, १०० र १५० के. जी. नाइट्रोजन प्रति हे.) र सहायक प्लटहरूमा गहुँको ६ वटा जातहरू (बी. एल. ३६२३, बी. एल. ३६२९, बी. एल. ३८७२, एन. एल. १००८, एन. एल. १०५५ र विजय) छरिएको थियो। दानाको उत्पादन र उत्पादनमा सघाउने आयामहरू परिक्षण गर्दा बढ्दो नाइट्रोजनको दुवै वर्षहरूमा समानान्तर असर देखियो। दुई वटा मापदण्डहरू पसाउने दिन र पाक्ने दिनहरू दुवै वर्षहरूमा जातहरू पिच्छे तथ्याङ्गीय रूपमा फरक देखिए। दुवै वर्षहरूमा सम्पूर्ण मापदण्डहरूमा गहुँका जातहरू र नाइट्रोजनको मात्राहरू बीच कुनै तथ्याङ्गीय अन्तरसम्बन्ध देखिएन। सन् २०१३ मा १०० के.जी. नाइट्रोजन प्रति हे. प्रयोग गर्दा विजय जातले ३.१२ टन प्रति हे. उत्पादन भएको पाइयो, भने सन् २०१४ मा १५० के.जी. नाइट्रोजन प्रति हे. को दरले प्रयोग गर्दा सोही जातको उत्पादकत्व ३.२३ टन प्रति हे. पाइयो। बालाको लम्बाई, प्रति वर्ग मि. मा रहेको उत्पादनमूलक गाँजहरूको संख्या, प्रति वर्ग मि. मा रहेको बालाहरूको संख्या र हजार दानाहरूको तौल बढ्दो नाइट्रोजनका मात्रा संगै बृद्धि भएको पाइयो। दुवै वर्षहरूमा १५० के. जी. नाइट्रोजन प्रति हे. प्रयोग गर्दा विजय जातले ४.३५ टन प्रति हे. र बी. एल. ३८७२ जातले ४.३३ टन प्रति हे. सबैभन्दा बढी परालको उत्पादन भएको पाइयो। सन् २०१३ मा विजय जातमा १०० के. जी. नाइट्रोजन प्रति हे. प्रयोग गर्दा सबैभन्दा बढी उत्पादनमूलक गाँजहरू २७६.३३ र सोही जातले सन् २०१४ मा १५० के.जी. नाइट्रोजन प्रति हे. को दरले प्रयोग गर्दा सबैभन्दा बढी उत्पादनमूलक २९७ गाँजहरू भएको पाइयो।

INTRODUCTION

Wheat is the third most important crop in terms of production in Nepal (Joshi 2015, Bhattarai et al 2015). Its area, production and yield were 7,54,243 ha, 17,27,346 t and 2.29 t ha⁻¹, respectively (MoAD 2013). In Sunsari district located in eastern Nepal, the productivity of wheat was 2.6 t ha⁻¹ in 2013 (DADO 2012/13). Since the wheat plays major role in human food and nutrition, thus after the entry of semi-dwarf wheat varieties, the trend of introducing advanced breeding lines has increased in Nepal. Despite of that, the production of wheat remained stagnant with a clear yield gap between national average and farmers' field (MoAD 2012/13). In addition, high variation could be observed among wheat genotypes with respect to phenological traits and responses to fertilizer application. The stagnant yield and gap could be caused by several reasons (NWRP 2012). One primary reason could be inappropriate and inadequate fertilization- thus, it is necessary to know the response of fertilizers among wheat genotypes for increasing production. Nitrogen (N) is generally the most common major limiting nutrient for growth and yield of crops worldwide (Camara et al 2003). The supply of nutrients in a suitable form is also a major determining factor for cereal production with N being the most important applied fertilizer (Spiertz 1984), as wheat is very sensitive to insufficient N and very responsive to N fertilization. N is one of the most important agronomic factors affecting the yield and technological quality, especially protein content, gluten amount and quality (Podolska 2014).

It should be noted that both an optimized N management for a less responsive cultivar and a restrictive management for a more demanding cultivar may result in crops with little yield potential (Russell et al 2014). High nutrient levels can also harm crops by making wheat plants more vulnerable to lodging (Ma et al 2010) and economic losses to farmers, because only 33% of all N fertilizers applied to cereal crops are absorbed in harvested grains (Raun and Johnson 1999). Thus, the use of N in wheat crops must be optimized to increase yields. There are reports of N use in wheat crops, ranging from 90 to 225 kg ha⁻¹ of N, without significant responses in grain yield under more favorable environment and management conditions (Barracough et al 2010). For irrigated wheat, Heinemann et al (2006) observed a positive response up to 156 kg ha⁻¹ of N, with a grain yield of 6472 kg ha⁻¹. So undoubtedly, N deficiency constitutes one of the major yield limiting factors for cereal production (Sarwar et al 2012).

Numerous researchers found that more N was demanded by modern wheat varieties, which is typically supplied in the form of N fertilizers (Campbell et al 1993, Wagen et al 2003 and Salvagiotti and Miralles 2008). The effectiveness of N depends on many factors

mainly the term of N applications and the variety (Podolska 2014). There are several studies on the use of N fertilizers and wheat production in Nepal (Rai and Khadka 2009, NWRP 2012, Bhattari et al 2015). However, in present work we elucidate the response of wheat genotypes to different levels of N fertilizers applied for the selection of most promising genotypes.

MATERIALS AND METHODS

Two field experiments were conducted during November-April of 2012/13 and 2013/14 on a clay loam soil at RARS, Tarahara (Figure 1). The GPS coordinates of experimental site is 26.7069 °N latitude and 087.2813 °E longitudes lying at 127 masl. The experiment was laid out in a split plot design with four nitrogen levels (0, 50, 100 and 150 kg N ha⁻¹) as main plots and six wheat genotypes (BL 3623, BL 3629, BL 3872, NL 1008, NL 1055 and Vijay) as sub-plots. Every treatment was replicated three times for a total of seventy two plots of 2 m x 5 m = (10 m²).

All other cultivation practices were uniform for all treatments. Composite soil samples from each block of the experiment were taken from the upper 25 cm depth before sowing. Land was thoroughly ploughed to make the soil well pulverized. Seed sowing was done on 30 November in both the years. Spacing was maintained 25 cm between rows and plot consisted of eight continuous rows of 5 m long. N was provided through urea and Diammonium phosphate (DAP). Phosphorus (P) was applied through DAP, but in no-nitrogen plot, the source of P was Single Super Phosphate and the source of potassium was Muriate of potash (MOP). Whole dose of P and potassium (K) were applied at the time of land preparation. Half N was applied in line at the time of sowing and the remaining half was broadcasted with the first irrigation at 25 DAS (days after sowing).



Figure 1. Location of the experimental site at RARS, Tarahara, Sunsari, Nepal.

The parameters recorded were (Table 1); emergence count m⁻², days to heading, days to maturity, plant height (cm), productive tillers m⁻², spike length (cm), number of spikes m⁻², test weight (g), grain yield (t ha⁻¹) and straw yield (t ha⁻¹). Grain and straw was brought from whole 10 m² for the measurement. Straw was sun dried for a day and weighed while grain weight was estimated at 14 % moisture. The measured weight of grain and straw were converted into hectare basis. The meteorological data was obtained from weather station that is supervised by Dharan based Department of Hydrology and Meteorology, Government of Nepal established at RARS, Tarahara. The methods adopted for the laboratory analysis has been given in Table 1. The soil characteristic of the experimental sites has been presented in Table 2. Data collected during the experiment were analyzed using MSTAT statistical package. The Duncan's multiple range (DMR) test at 5% probability level was used to test the differences among mean values.

Table 1. Parameters and methods adopted for the laboratory analysis at RARS, Tarahara

SN	Parameter	Method
1	pH	Potentiometric 1:2 (Jackson 1973)
2	Soil Organic Matter (%)	Walkely and Black (Walkely and Black 1934)
3	Total N (%)	Kjeldahl (Bremner and Mulvaney 1982)
4	Available P (kg ha ⁻¹)	Olsen (Olsen et al 1954)
5	Available K (kg ha ⁻¹)	Ammonium acetate (Jackson 1967)
6	Soil texture (Particle size analysis)	Hydrometer (Bouyoucos 1927)

Table 2. Soil characteristics of experimental site at RARS Tarahara, Nepal

Characteristics	2012	2014
pH	7.2	6.5
Soil organic matter (%)	4.1	3.7
Total N (%)	0.12	0.17
Available P (kg ha ⁻¹)	119	115
Available K (kg ha ⁻¹)	136	119
Soil texture (Particle size analysis)	Clay loam	Clay loam




Figure 2. Meteorological data for the year 2013 (A) and 2014 (B) at RARS, Tarahara, Nepal.

RESULTS

The meteorological data of the study area has been presented in Figure 2. Mainly three parameters were monitored, rainfall pattern, minimum and maximum air temperature (Figure 2). The pattern of rainfall was almost similar in both the years, however, in 2014 there was slightly earlier rainfall even in the month of January, comparative to 2013 the rainfall in 2014 was slightly higher in the months of March and April, while the pattern of minimum and maximum air temperature was almost similar in both years lowest in January and February in 2014. N application significantly ($p \leq 0.05$) affected all parameters tested in both the years (Table 3). Of all the parameters, days to heading (DTH) and days to maturity (DTM) significantly ($p \leq 0.05$) varied among six wheat genotypes in both the years. In both years, BL 3872 took the maximum days to heading ranging from 87.92 to 89.82 days. The maximum days to maturity (120) was observed in NL 1055 in 2013, while BL 3872 took the maximum days to maturity (126.5) in 2014. The effect of interaction between N and wheat genotypes on all parameters during both the years was not significant ($p \leq 0.05$).

Table 3. Growth, yield and yield components of wheat genotypes under different levels of nitrogen in 2013 at RARS, Tarahara, Nepal

Treatments	Emergence count m^{-2}	Days to heading	Days to maturity	Plant height (cm)	Productive tillers m^{-2}	Spike length (cm)	Spikes m^{-2}	Test weight (g)	Grain yield ($t ha^{-1}$)	Straw yield ($t ha^{-1}$)
N levels										
N ₀	130.6 ^c	79.5 ^c	107.7 ^c	77.74 ^c	143.6 ^c	8.33 ^b	149.4 ^d	38.64 ^d	1.67 ^c	2.28 ^c
N ₁	141.4 ^b	84.44 ^b	115.2 ^b	84.68 ^b	164.6 ^b	9.46 ^a	167.8 ^c	41.20 ^c	2.30 ^b	2.65 ^c
N ₂	148.9 ^b	88.44 ^a	116.6 ^b	88.81 ^a	251.8 ^a	9.66 ^a	201.1 ^b	44.28 ^b	2.88 ^a	3.56 ^b
N ₃	163.2 ^a	90.61 ^a	124.3 ^a	90.22 ^a	264.9 ^a	10.32 ^a	220.0 ^a	46.40 ^a	2.90 ^a	4.19 ^a
F test	**	**	**	**	**	**	**	**	**	**
LSD (0.05)	10.23	3.79	2.89	1.94	19.4	0.89	17.28	1.67	0.280	4.47
SEm (±)	157.3	21.5	12.49	5.64	565.9	1.19	449.0	4.21	0.12	0.33
Genotypes										
BL 3623 (V1)	145.8	83.67 ^b	113.4 ^{bc}	85.65	208.5	9.11	174.1	44.23	2.47	3.24
BL 3629 (V2)	148.1	83.5 ^b	115.5 ^{abc}	85.14	202.2	9.37	186.3	42.16	2.41	2.96
BL 3872 (V3)	146.6	87.92 ^a	118.2 ^{ab}	82.52	204.3	9.41	190.3	42.83	2.48	3.18
NL 1008 (V4)	144.0	87.83 ^a	116.4 ^{abc}	87.14	207.8	9.72	175.8	42.07	2.33	3.25
NL 1055 (V5)	148.1	85.75 ^{ab}	120.0 ^a	86.75	209.2	9.31	186.8	42.13	2.40	3.14
Vijay (V6)	143.8	85.83 ^{ab}	112.3 ^c	87.02	213.0	9.75	194.3	45.37	2.52	3.26
F test	NS	*	*	NS	NS	NS	NS	NS	NS	NS
LSD (0.05)	-	2.50	5.80	-	-	-	-	-	-	-
SEm (±)	107.1	9.17	49.47	3.07	67.16	0.44	355	7.74	0.43	0.15
CV %	7.09	3.53	6.67	7.92	8.47	7.04	10.21	6.52	8.53	11.99
Grand mean	146.04	85.75	115.96	86.37	207.49	9.44	184.58	42.63	2.44	3.17
Interaction										
NxV	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

Emergence count m^{-2} of wheat genotypes had significantly ($p \leq 0.05$) affected in both the years by different levels of N applied. Results indicated that the highest emergence m^{-2} , 163.2 in the year 2013 and 164.8 in the year 2014 were recorded when the crop was fertilized with 150 kg N ha^{-1} (Table 3 and 4). In both the years days to heading (DTH) and days to maturity (DTM) were significantly ($p \leq 0.05$) affected by different N levels (Table 3 and 4). The maximum plant height (90.22 cm) was resulted in treatment N3 (150 kg N ha^{-1}), but was statistically ($p \leq 0.05$) similar (88.81 cm) to N2 (100 kg N ha^{-1}) in 2013.

In 2014, the application of 150 kg N ha^{-1} gave also the highest plant height (92.43 cm) in same treatment. However, average plant height was lower in the year 2014 (85.30) as compared to previous year (86.37 cm) (Table 3 and 4). The N treatments significantly ($p \leq 0.05$) increased the numbers of productive tillers m^{-2} than in control in both the years. Application of 150 kg N ha^{-1} produced the maximum numbers of productive tillers (264.9 m^{-2}) and (273.6 m^{-2}) in 2013 and 2014, respectively. The check variety, Vijay produced the maximum numbers of productive tillers in both the years with 213 m^{-2} and 224.6 m^{-2} . This trend was attributed to comparatively better growth due to higher fertilizer application. Spike length increased with increasing level of N; 100 kg N ha^{-1} and 150 kg N ha^{-1} produced statistically similar result in 2013 (Table 3 and 4). However, application of 150 kg N ha^{-1} resulted the maximum spike length (11.24 cm) followed by 10.09 cm in case of N₃ (100 kg N ha^{-1}) in 2014. The maximum number of spikes m^{-2} was produced by the application of 150 kg N ha^{-1} in both the years i.e. 220 m^{-2} and 249 m^{-2} in 2013 and 2014, respectively.

The highest values of 1000-grain weight 46.40 g and 46.90 g were achieved in 2013 and 2014 respectively, both with the application of 150 kg N ha^{-1} . Hence, the yield contributing characters viz. productive tillers m^{-2} , spike length (cm), spikes m^{-2} and test weight (g) increase significantly ($p \leq 0.05$) with every increase in the dose of N up to 150 kg N ha^{-1} in both the years (Table 3 and 4). Grain yield response varied with the season and yield was higher in 2014 seasons than 2013. The average increment of grain yield in 2014 compared to 2013 was 5.52% at 150 kg N ha^{-1} . In both the years, Vijay resulted the highest grain yield; 3.12 t ha^{-1} in 2013 with the application of 100 kg N ha^{-1} and 3.23 t ha^{-1} in 2014 by 150 kg N ha^{-1} followed by BL 3872 in 2013 (3.01 t ha^{-1}) and BL 3623 in 2014 (3.16 t ha^{-1}), both with the application of 150 kg N ha^{-1} (Table 3 and 4). In both the years, N₂ (100 kg N ha^{-1}) and N₃ (150 kg N ha^{-1}) produced statistically similar results (Table 3 and 4). The obtained grain yield in wheat might be the result of effective tillers, number of spikes m^{-2} and grain weight. The highest straw yield (4.19 t ha^{-1}) was obtained from application of 150 kg N ha^{-1} followed by 3.56 t ha^{-1} in case of 100 kg N ha^{-1} in 2013. Similarly, the maximum straw yield (3.66 t ha^{-1}) was obtained from 150 kg N ha^{-1} in 2014 (Table 4).

Table 4. Growth, yield and yield components of wheat genotypes under different levels of nitrogen in the year 2014 at RARS, Tarahara, Nepal

Treatments	Emergence count m^{-2}	Days to heading	Days to maturity	Plant height (cm)	Productive tillers m^{-2}	Spike length (cm)	Spikes m^{-2}	Test weight (g)	Grain yield (t ha^{-1})	Straw yield (t ha^{-1})
N levels										
N ₀	137.8 ^c	81.17 ^b	108.4 ^d	79.53 ^c	146.6 ^d	8.370 ^c	151.7 ^c	39.16 ^d	1.75 ^c	2.56 ^c
N ₁	147.6 ^{bc}	82.61 ^b	116.5 ^c	85.93 ^{bc}	165.5 ^c	9.479 ^b	172.6 ^c	41.85 ^c	2.53 ^b	2.74 ^c
N ₂	159.0 ^{ab}	86.89 ^a	121.1 ^b	86.61 ^b	242.9 ^b	10.09 ^b	215.1 ^b	44.79 ^b	2.94 ^a	3.38 ^b
N ₃	164.8 ^a	89.83 ^a	131.2 ^a	92.43 ^a	273.6 ^a	11.24 ^a	249.9 ^a	46.90 ^a	3.06 ^a	3.66 ^a
F test	**	**	**	**	**	**	**	**	**	**
LSD _(0.05)	12.35	3.81	2.88	14.16	16.81	0.63	30.34	1.19	0.13	0.28
SEm (±)	229.4	21.86	12.49	301.40	425.0	0.59	1383	2.14	0.03	0.12
Genotypes										
BL 3623 (V1)	156.2	82.75 ^b	118.2 ^{bc}	84.46	209.6	9.52	178.6	44.28	2.60	3.01
BL 3629 (V2)	146.3	83.08 ^b	113.6 ^c	83.68	201.0	9.56	200.3	42.89	2.64	2.84
BL 3872 (V3)	147.7	89.82 ^a	126.5 ^a	86.29	204.6	9.90	197.8	42.67	2.52	3.34
NL 1008 (V4)	150.6	85.75 ^{ab}	118.2 ^{bc}	85.16	206.2	9.99	197.1	43.17	2.49	3.15
NL 1055 (V5)	155.9	84.67 ^b	120.8 ^{ab}	85.39	198.8	9.76	184.3	44.19	2.42	3.28
Vijay (V6)	155.6	85.58 ^{ab}	118.6 ^{bc}	86.84	224.6	10.03	212.7	42.72	2.74	3.08
F test	NS	*	*	NS	NS	NS	NS	NS	NS	NS
LSD _(0.05)	-	3.66	6.49	-	-	-	-	-	-	-
SEm (±)	85.13	19.67	61.77	9.17	464.7	0.48	687.8	9.47	0.09	0.13
CV %	6.07	5.21	6.59	5.02	10.41	7.06	13.29	7.13	11.47	12.32
Grand mean	152.03	85.13	119.29	85.30	207.125	9.79	197.33	43.18	2.59	3.09
Interaction										
NxV	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

DISCUSSION

The present data were supportive for the evaluation of nutritional demand and yield potential of different pipeline wheat varieties for N fertilization. It was shown that higher performance of yield components and ultimate grain yield was associated with higher N fertilization levels (Table 4). The significant ($P \leq 0.05$) enhancement of plant height in all treatments over the control (N_0) in both the years can be credited to higher dose of N, which greatly helps the plant to expose its potential to grow potentially. The increasing dose of N result the linear increment in spike length in both years for all pipeline varieties. These results are in conformity with those of (Rai and Khadka 2009) who observed maximum spike length under fertilization of 100 kg N ha^{-1} as compared to control and 50 kg N ha^{-1} in their long term experiment in Khumaltar. Higher test weight with increasing N levels might be due to supply of more N and higher production of photosynthates (Naseri et al 2010) that showed linearly increased test weight due to increasing levels of N application (Figure 4).

Hence, the yield contributing characters viz productive tillers m^{-2} , spike length (cm), spikes m^{-2} and test weight (g) increase significantly ($p \leq 0.05$) with every increase in the dose of N up to 150 kg N ha^{-1} in both the years (Table 3 and 4). This tendency was attributed to comparatively better growth due to higher fertilizer application. The poor growth of the wheat plants and the consequential lower grain yield in control (N_0) plots may be due to competition for N from the soil and starved conditions. Meteorological conditions for the two years were more suitable to the emergence and survival of tillers and further tillering and development of spikes at higher nitrogen levels (Figure 2 and 3).




Figure 3. Grain yield of wheat genotypes from different treatments in years 2013 and 2014.

Probably the meteorological suitability enhanced competition for nutrients and photo assimilates making the varieties more responsive to the applied fertilizers. A greater response to N fertilization is obvious when a meteorological condition remains favorable, especially rainfall were not the limiting factors (Figure 2 A, B). The responsiveness of the tested varieties to the applied fertilizers should be studied further before it reached to farmers' field.

Benis et al (1994) argued that the yield response of wheat to nitrogen depends on factors such as climate, soil and cultivar. Identification of correlations between meteorological variables and grain yield for such pipelines can be another subject of study.




Figure 4. Grain yield of wheat genotypes to various N levels in 2013 (A) and 2014 (B).

CONCLUSION

The study showed that all the wheat genotypes responded well up to the levels of 150 kg N ha⁻¹ regarding yield and yield attributing characters. The performance of Vijay with respect to productive tillers m⁻², grain yield and straw yield with combination of 150 kg N ha⁻¹ was the best. With well cultural management practices application of 100 kg N ha⁻¹ can give good grain yield of wheat.

ACKNOWLEDGEMENT

The authors express their gratitude to the government of Nepal, Nepal Agricultural Research Council (NARC) for supporting this study. This is an outcome of Project 40370004 funded by NARC. The authors are thankful to RARS (Tarhara) and National Wheat Research Program, Bhairahawa, Rupandehi for providing seeds, accessories and other supports.

REFERENCES

- Barraclough PB, JR Howarth, J Jones, R Lopez-Bellido, S Parmar, CE Shepherd and MJ Hawkesford. 2010. Nitrogen efficiency of wheat: genotypic and environmental variation and prospects for improvement. *European Journal of Agronomy* 33:1-11.
- Benin G, E Bornhofen, E Beche, E Stefani Pagliosa, C Lemes da Silva and C Pinnow. 2012. Agronomic performance of wheat cultivars in response to nitrogen fertilization levels. *Acta Scientiarum* 34(3):275-283.
- Bhattarai R, B Chaudhary, DB Thapa, RR Puri, RN Chaudhary, B Sapkota, KB, SR Shrestha and SP Adhikari. 2015. Wheat yield interaction with sowing dates of Seeding in Eastern Mid Hills of Nepal, *Journal of Nepal Agricultural Research Council*, 1:33-36.
- Bouyoucos GJ. 1962. Hydrometer method improved for making particle size analyses of soils. *Agronomy Journal* 54(5):464-465.
- Bremner J M and CS Mulvaney. 1982. Nitrogen total. In: A. L. Page (ed.), *Methods of soil analysis*. Agron. No. 9, Part 2: Chemical and microbiological properties. 2nd edition. Am. Soc. Agron., Madison, W I, USA. pp.595-624.
- Camara KM, WA Payne and PE Rasmussen. 2003. Long term effects of tillage, nitrogen, and rainfall on winter wheat yields in the Pacific Northwest. *Agron. J.* 95:828-835.
- Campbell CA, RP Zenner, F Selles, BG McConkey and FB Dyck. 1993. Nitrogen management for spring wheat grown annually on zero tillage: yields and nitrogen use efficiency. *Agron. J.* 85(1):107-114.
- DADO 2012/13. Annual Agriculture Development Program and Statistical Book. District Agriculture Development Office, Inaruwa, Sunsari. 67
- Heineman A B, LF Stone, DD Agostinho and AD Canovas. 2006. Solar radiation use efficiency on the wheat grain yield as a function of nitrogen fertilizer. *Rev. bras. Eng. Agric. Ambient* 10:352-356.
- Jackson M L. 1967. Handling soil samples in the laboratory. In: *Soil chemical analysis*. Prentice hall of India, Pvt. Ltd., New Delhi 2: 30-37.
- Jackson ML. 1973. *Soil Chemical Analysis*. Pentice Hall of India Pvt. Ltd., New Delhi.
- Joshi BK, Thapa DB, Bhatta MR .2015. Genetic parameters of common wheat, *Journal of Nepal Agricultural Research Council*,1:9-13.
- Ma BL, TY Wu, N Tremblay, W Deen, NB McLaughlin, MJ Morrison and G Stewart. 2010. On-farm assessment of the amount and timing of nitrogen fertilizer on ammonia volatilization. *Agronomy Journal* 102:134-144.
- MoAD. 2012/13. Crop situation update. A joint assessment of 2012/13 winter and 2013/14 summer crops outlook, MoAD, Nepal. 18pp.
- MoAD. 2013. Statistical Information on Nepalese Agriculture. 2013. Agribusiness Promotion and Statistics Division, Statistics Section, Ministry of Agricultural Development, Government of Nepal, Singh Durbar, Kathmandu, Nepal.
- Naseri R, A Mirzaei, R Soleimani and E Nazarbeygi. 2010. Response of bread wheat to nitrogen application in calcareous soils of Western Iran. *Am-Eurasian J. of Agric. Env. Sci.* 9(1):79-85.
- NWRP. 2012. Executive summary. In: Annual report-2012. National Wheat Research Program (NWRP)- Nepal Agricultural Research Council, Bhairahawa- Rupandehi. pp. 6-11.
- Olsen SR, CV Cole and LA Dean. 1954. Estimation of available phosphorus in soil by extraction with sodium carbonate. In: *Method of soil analysis, Part 2*. (CA Black, ed) American Soc. Agron. Inc, Medison, USA. pp. 1044-1046.
- Podolska G. 2014. Technological quality of winter wheat depending on nitrogen applications. In: *Proceedings of the 18th Nitrogen Workshop- The nitrogen challenge: building a blueprint for nitrogen use efficiency and food security* (CMdS Cordovil, ed). 30th June-3rd July 2014, Lisbon, Portugal. 175 p.
- Rai S K and Y G Khadka. 2009. Wheat production under long term application of inorganic and organic fertilizers in rice-wheat system under rainfed conditions. *Nepal Agric. Res. J.* 9:123-131.
- Raun WR and GV Johnson. 1999. Improving nitrogen use efficiency for cereal production. *Agronomy Journal* 91:357-363.
- Riley WJ, I Ortiz-Monasterio and PA Matson. 2001. Nitrogen leaching and soil nitrate, nitrite, and Wheat responsiveness to nitrogen. *Nutrient Cycling in Agroecosystems* 61:223-236.
- Rusell K, DV Sanford and C Lee. 2014. Genotype specific management for nitrogen use efficiency in Kentucky soft red winter wheat. *Aspects of Applied Biology* 25:227-287.
- Salvagiotti F and DJ Miralles. 2008. Radiation interception, biomass production and grain yield as affected by the interaction of nitrogen and sulfur fertilization in wheat. *Europ. J. Agron.* 28:282-290.
- Sarwar M, G Jilani, E Rafique, ME Akhtar and AN Chaudhry. 2012. Impact of integrated nutrient management on yield and nutrient uptake by maize under rain-fed conditions. *Pak. J. Nutr.* 11:27-33.
- Spiertz J, D Vos and L Hole. 1984. The role of nitrogen in the yield formation of cereals, especially of winter wheat. In: *the Proceedings of Cereal Production*, Royal Dublin Society, Bultworths 5(3):34-65.
- Wagen M R, FC Oad and KS Nenwani. 2003. Quantitative and qualitative characteristics of wheat crop under various sources and methods of nitrogenous fertilizer application. *Asian J. of Plant. Sci.* 2(9):683-687.
- Walkley AJ and IA Black. 1934. Estimation of soil organic carbon by the chromic acid titration method. *Soil Sci.* 37(1):29-38.