



Original Article

Prevalence of Helicobacter pylori among patients with dyspepsia and correlation between endoscopic and histological diagnosis.

KC Shiva Raj¹, Lakhey A¹, Koirala K², Amatya GL³

¹Department of Pathology, KIST Medical College, Lalitpur, Nepal.

²Department of Surgery, KIST Medical College, Lalitpur, Nepal.

³Department of Gastroenterology, Bir Hospital, Kathmandu, Nepal.

Keywords:

Gastritis;
Follicular gastritis;
Carcinoma;
Endoscopy;
Active gastritis

ABSTRACT

Background: Dyspepsia is a prevalent complaint in general practice and gastrointestinal clinics. Helicobacter pylori have major causal relationship with gastro duodenal disease. The following study seeks to identify the prevalence of H. pylori based on histology and to correlate endoscopic findings with histopathology.

Materials and Methods: This was a cross-sectional observational study conducted in GRP Polyclinic and Om Hospital and research centre from April 2015-September 2015. The upper gastrointestinal endoscopic findings were recorded and were correlated with histopathological findings. All the relevant data were collected and analysed using Statistical Package of Social Sciences version 16 for windows.

Results: Endoscopy finding was divided into reflux esophagitis, antral gastritis, duodenitis, duodenal ulcer, gastric ulcer, and gastric cancer. Duodenal ulcer and gastric ulcer was noted more frequently in males than in females (55.0% vs. 45.0% and 58.2% vs. 41.8%), respectively, $P < 0.001$.

Chronic follicular gastritis was the most common in gastric ulcer (41.7%), whereas chronic persistent gastritis was common in non-ulcerative disease. Chronic active gastritis and chronic follicular gastritis were more common in ulcerative diseases, whereas chronic persistent gastritis was more common in gastritis and duodenitis ($P < 0.001$). The overall prevalence of H. pylori infection was 68.1% with male preponderance. Chronic active gastritis had highest prevalence of H. pylori (84.8%), followed by chronic follicular gastritis (84.1%) and chronic persistent gastritis (p value < 0.001).

Conclusion: Rate of H. pylori infected patients with dyspepsia was high. Ulcerative lesions were more common in males than in females with higher rate of infection with H. Pylori. Histological diagnosis of chronic active gastritis and chronic follicular gastritis was the most common pathologies in ulcerative lesions.

INTRODUCTION

Dyspepsia is a prevalent complaint in general practice and gastrointestinal clinics.¹ Helicobacter pylori (H. pylori) were of major concern today because of its causal relationship with gastro duodenal disease. After the discovery of H. pylori by Marshall and Warren in 1983 by using Warthin Starry Silver stain, the etiological understanding of gastritis

Correspondence:

Dr. Shiva Raj K C, MBBS, MD

Department of Pathology

KIST Medical College, Imadole, Lalitpur, Nepal

Email: shiva_kc_123@yahoo.com

Table: 1 Prevalence of Helicobacter Pylori in accordance with endoscopic diagnosis

Endoscopy	Male no (%)	Female no (%)	Overall no (%)
Esophagitis	20/35(57.0)	30/67 (44.7)	50/102 (49.0)
Gastritis	151/246 (61.3)	222/337 (65.8)	373/583 (63.9)
Duodenitis	49/73 (67.1)	27/34 (79.4)	76/107 (71.0)
Gastric Ulcer	56/60 (93.3)	34/43 (79.0)	90/103 (87.3)
Duodenal ulcer	57/66 (86.3)	45/54 (83.3)	102/120(85.0)
Gastric carcinoma	3/3(100)	1/2 (50)	4/5 (80)
Total	336/483 (69.5)	359/537 (66.8)	695/1020 (68.1)

P=0.043

Table: 2 Incidence of Helicobacter Pylori in accordance with endoscopic diagnosis and correlation with histopathology

Diagnosis	Chronic persistent Gastritis			Chronic Active Gastritis			Chronic Follicular Gastritis			Carcinoma		
	H.Pylori +	H.Pylori -	Total	H.Pylori +	H.Pylori -	Total	H.Pylori +	H.Pylori -	Total	H.Pylori +	H.Pylori -	Total
Esophagitis	16/41 (39.0%)	25/41 (61.0%)	41/102 (40.19%)	30/56 (53.57%)	26/56 (46.42%)	56/102 (54.90%)	4/5 (80%)	1/5 (20%)	5/102 (4.90%)	-	-	-
Gastritis	93/251 (37%)	158/251 (62.95%)	251/583 (43%)	190/225 (84.4%)	35/225 (15.5%)	225/583 (38.6%)	90/107 (84.1%)	17/107 (15.8%)	107/583 (18.3%)	-	-	-
Duodenitis	22/53 (41.5%)	31/53 (58.5%)	53/107 (49.5%)	40/40 (100%)	0/40 (0.0%)	40/107 (37.3%)	14/14 (100%)	0/14 (0.0%)	14/107 (13.0%)	-	-	-
Duodenal Ulcer	13/27 (48.1%)	14/27 (51.8%)	27/120 (22.5%)	67/67 (100%)	0/67 (0.0%)	67/120 (62.6%)	22/26 (84.6%)	4/26 (15.3%)	26/120 (21.6%)	-	-	-
Gastric Ulcer	18/18 (100%)	0/18 (0.0%)	18/103 (17.4%)	38/42 (90.4%)	4/42 (9.5%)	42/103 (40.7%)	34/43 (79.0%)	9/43 (20.9%)	43/103 (41.7%)	-	-	-
Carcinoma	-	-	-	-	-	-	-	-	-	4/5 (80.0%)	1/5 (20.0%)	5/5 (100.0%)
Total	162/390 (41.5%)	228/390 (58.4%)	390/1020 (38.2%)	365/430 (84.8%)	65/430 (15.1%)	430/1020 (42.1%)	164/195 (84.1%)	31/195 (15.9%)	195/1020 (19.1%)	4/5 (80.0%)	1/5 (20.0%)	5/5 (100.0%)

and its role as bacterial carcinogen have changed the management of gastritis. One half of the world's population has H. pylori infection, with an estimated prevalence of more than 90% in developing countries.²⁻⁶ In our country, the reported prevalence of H. pylori ranged from 30% to 67%.⁷⁻¹¹ This bacterium colonizes human gastric mucosa and can elicit lifelong inflammatory and immune responses, with release of various bacterial and host dependent cytotoxic substances. It causes chronic and active gastritis, peptic ulcer disease and associated with increased risk of developing gastric cancer.¹²⁻¹⁴

The following study seeks to identify the prevalence of H. pylori based on histology and to correlate endoscopic findings with histopathology.

MATERIAL AND METHODS

This was an observational cross-sectional study at GRP Polyclinic Pvt. Ltd and Om Hospital and Research centre Pvt Ltd over a period of 6 months from from April 2015-September 2015. Data analysis was done from all consecutive individuals who had undergone gastric biopsy during upper GI endoscopy for various dyspeptic symptoms like pain abdomen, nausea, vomiting, belching, throat pain, upper gastrointestinal bleeding, weight loss etc. However, patients under the age of 14 years, with oesophageal varices were not included in the study. Prior to investigation, informed consent was taken from the patient. Permission

from ethical review committee was obtained.

The upper gastrointestinal endoscopy was performed by two physicians separately and endoscopic findings were categorized into gastro-esophageal reflux disease, antral gastritis, gastric ulcer, duodenitis, duodenal ulcer, and neoplasm. The biopsy specimens were usually taken from the gastroesophageal junction, antrum or duodenum depending upon the endoscopic findings and sent for histological examination. The biopsy specimens were fixed in 10% buffered formalin, processed, embedded in paraffin, and cut and stained with Hematoxylin and Eosin (HandE) and Giemsa stain. Histological reporting was done by pathologist using Modified Sidney system which included inflammation, activity, atrophy, intestinal metaplasia, and Helicobacter Pylori colonization.¹⁵ Histologically, gastritis was classified into chronic active gastritis (CAG), chronic follicular gastritis (CFG), and chronic persistent gastritis (CPG) with or without intestinal metaplasia, atrophy and Helicobacter Pylori colonization.

All the relevant data were collected and analysed using Statistical Package of Social Sciences (SPSS) version 16 for windows. Chi-square test with exact test was used where applicable. P values of < 0.05 were considered to denote statistical significance.

RESULTS

TA total of 1020 UGI endoscopies were performed and biopsies were taken during the study period. The age of the patients ranged from 16-94 years with the mean of 41.7 ±18. Among all the patients, 549 (53.8%) were females and 471 (46.1%) were males.

Endoscopy finding was divided into reflux esophagitis, antral gastritis, duodenitis, duodenal ulcer, gastric ulcer, and gastric cancer, which was diagnosed in 102 (10.0%), 583 (57.1%), 107 (10.5%), 120 (11.7%), 103 (10.0%), and 5 (0.5%) patients, respectively. Duodenal ulcer and gastric ulcer was noted more frequently in males than in females (n = 66; 55.0% vs. n = 54; 45.0% and n = 60; 58.2% vs. n = 43; 41.8%), respectively, P < 0.001). In contrast, Esophagitis, antral gastritis was more common among females (n = 67; 65.6% vs. n=35; 34.4% and n=337; 57.8% vs. n = 246; 42.2%) respectively.

The histological features were divided into chronic persistent gastritis (CPG), chronic active gastritis (CAG), chronic follicular gastritis (CFG), and gastric carcinoma which were detected in 390 (38.2%), 430 (42.1%), 195 (19.1%), and 5 (0.5%) of cases respectively. CAG was the frequent finding in those who had duodenal ulcer (n = 67; 62.6%), all of which (100%) were colonized with *H. pylori* followed by gastric ulcer (n = 42; 40.7%). CFG was most common in gastric ulcer (n = 43; 41.7%), whereas CPG was common in non-ulcerative disease (duodenitis - n = 53; 49.5%; and gastritis - n = 251; 43%). CAG and CFG were more common in ulcerative diseases, whereas CPG was more common in gastritis and duodenitis (P < 0.001). Those who had histological features of CAG had highest prevalence of *H. pylori* (n = 365; 84.8%), followed by CFG (n = 164; 84.1%) and CPG (n = 162; 41.5%) with P value < 0.001.

The overall prevalence of *H. pylori* infection was 695/1020 (68.1%) with male preponderance; Male: 69.5% vs. F: 66.8% (Table 1). *Helicobacter Pylori* was found in 50 (49.0%) cases of esophagitis, 373 (63.9%) cases of antral gastritis, 76 cases (71.0%) of duodenitis, 102 (85.0%) cases of duodenal ulcer, 90 cases (87.3%) of gastric ulcer, and 4 cases (80%) of gastric carcinoma. In gastric ulcer and duodenal ulcer including gastric carcinoma there was significantly higher prevalence of *H. pylori* (P = 0.046); however, no significant difference in the prevalence of *H. pylori* was observed between esophagitis, gastritis and duodenitis. The colonization rate of *H. pylori* in relation to different endoscopy diagnosis is illustrated in Table 1.

Intestinal metaplasia was identified in 3% (31/1020) of the patients who underwent UGI endoscopy. Only 11 (35.4%) patients with intestinal metaplasia had colonization by *Helicobacter Pylori*. The mean age of the patient with intestinal metaplasia was 54.2 ± 16.9 years. There was

significant male predominance with male to female ratio of 1.6:1.

Atrophy was noted in 24(2.35%) of all the patients included in this study. There was female predominance male to female ratio of 0.7:1. The mean age of the patient with atrophic gastritis was 40.8±12.6.

DISCUSSION

Current standard of care for most patients with symptoms of upper GIT consists of initial endoscopic examination followed by histopathology. The individual accuracy and sensitivity of these techniques in the diagnosis of any lesions is subjective to the operator's ability to target site of pathology. *Helicobacter Pylori* remains the most common cause of chronic gastritis. Furthermore, since *Helicobacter pylorus* has carcinogenic effect, it has radically changed our understanding and clinical management of gastroduodenal disease, and much has been researched about its clinical aspects and its epidemiology.¹⁶

In this study age of the patients with dyspepsia ranged from 16-94 years with the mean of 41.7 years. Among the patients seeking for medical help due to dyspepsia 549 (53.8%) were female and 471(46.1%) were male with slight female predominance (M:F- 1:1.16). A study done by Sharma et al documented 51% female seeking for medical help due to dyspepsia as compared to 49% males.¹⁷

Most common upper gastrointestinal endoscopic findings were antral gastritis (57.1%) followed by duodenal ulcer (11.7%), duodenitis (10.5%) and gastric ulcer (10.0%) Similar finding was observed in a study done by Poudel A et al. He observed 58.1% cases had antral gastritis followed by 11.63% with peptic ulcer.¹⁸ Reflux esophagitis was observed in 10% of the patients which was less than study conducted by Ercelep OB et al.¹⁹ Ulcerative lesions of stomach and duodenum was more common in males than in females as was observed in a study done by Shrestha R et al.²⁰

Most common pathology in histopathology was chronic active gastritis (42.1%), followed by chronic persistent gastritis (38.2%), chronic follicular gastritis (19.1%) and gastric carcinoma (0.5%). This result was in concordance with other study.²⁰ In ulcerative lesions most common antral biopsy finding was chronic active gastritis and chronic follicular gastritis whereas in non-ulcerative lesions chronic persistent gastritis was the commonest diagnosis. Patients with chronic active gastritis and chronic follicular gastritis had higher frequency of *Helicobacter Pylori* positivity with incidence of 84.8% and 84.1% respectively. Higher incidence of *Helicobacter Pylori* were also seen in other studies done by Shrestha R et al.²⁰ Other studies also identified association between ulcerative lesions and higher incidence of *Helicobacter Pylori*.²¹⁻²³ Study done by X Y Chen et al observed positive correlation with the prevalence

and density of lymphoid follicles and aggregates with H pylori infection.²⁴ In a study done by Genta RM et al²⁵ observed 100% presence of Helicobacter Pylori in patients with chronic follicular gastritis. These discrepancies may be due to: 1) Various biopsy sites, 2) intake of proton pump inhibitor which leads to subepithelial colonization of bacteria and is difficult to identify in routine histopathology procedure. Overall prevalence of H. Pylori infection was 68.1%. Overall, researchers found a consistent pattern in most developing nations, where 70 to 90% of adults harbored the bacteria; most individuals acquired the infection as children, before age ten.^{5,6} Nepalese data show that the prevalence of H. pylori infection varies widely from 30-67%.⁶⁻¹¹

Intestinal metaplasia was observed in 3% (n=31) of the dyspeptic patients whereas gastric atrophy was noted in 3.25% (n=24) patients. Gastric carcinoma was seen in 5 (0.5%) of cases with 80% having Helicobacter Pylori infection. The data is too small to compare with other studies.

CONCLUSION

Rate of H. pylori infected patients with dyspepsia was high. Ulcerative lesions were more common in males than in females with higher rate of infection with H. Pylori. detected by histopathology among adult patients with gastrointestinal symptoms was high (68%). Histological diagnosis of chronic active gastritis and chronic follicular gastritis was the most common pathologies in ulcerative lesions. Because of higher prevalence rate of H. Pylori; routine endoscopic biopsy should be performed so as to detect this gram negative microorganism and to treat accordingly.

REFERENCES

1. Sobieraj, D.M., Coleman, S.M. and Coleman, C.I. (2011) US Prevalence of Upper Gastrointestinal Symptoms: A Systematic Literature Review. *American Journal of Managed Care* 2011;17:449-58.
2. Go MF. Natural history and epidemiology of Helicobacter pylori infection. *Aliment Pharmacol Ther* 2002;16:3-15. Crossref
3. Megraud F. Epidemiology of Helicobacter pylori infection. In: Rathbone BK, Heatley RV, editors. *Helicobacter pylori and gastroduodenal disease*. London: Blackwell Scientific Publication; 1993. pp.107-23.
4. Veldhuyzen van zantel SJ. Do socioeconomic status, marital status and occupation influence the prevalence of Helicobacter pylori infection? *Aliment Pharmacol Ther*.1995;9:41-4.
5. Hunt RH, Xiao SD, Megraud F, Leon-Barua R, et al. Helicobacter pylori in developing countries. *World Gastroenterology Organisation Global Guidelines; World Gastroenterology Organization. J Gastrointest Liver Dis*. 2011;20:299-304. Crossref
6. Lacy BE, Rosemore J. Helicobacter pylori: Ulcers and more: The beginning of an era. *J Nutr*.2001;131:27895-935. [PubMed]
7. Rai SK, Shah RD, Bhattachan CL, Rai CK, Ishiyama S, Kurokawa M, et al. Helicobacter pylori associated gastroduodenal problem among the Nepalese. *Nepal Med Coll J*. 2006;8:9-13. Crossref
8. Makaju R, Tamang MD, Sharma Y, et al. Comparative study on a homemade rapid urease test with gastric biopsy for diagnosis of Helicobacter pylori infection. *Nepal Med Coll J*. 2006;8:97-100. Crossref
9. Makaju RK, Tamang MD, Sharma Y, Sharma N, Koju R, Ashraf M. Prevalence of Helicobacter pylori in Dhulikhel Hospital, Kathmandu University Teaching Hospital: A retrospective histopathologic study. *Kathmandu Univ Med J* 2005;3:355-9.
10. Khanal S, Rao BS, Sharma Y, Khan GM, Makaju R. Comparative study between two triple therapy regimens on eradication of Helicobacter pylori. *Kathmandu Univ J Sci Engg Technol*. 2005:1-10.
11. Kawasaki M, Kawasaki T, Ogaki T, et al. Seroprevalence of Helicobacter pylori infection in Nepal: Low prevalence in an isolated rural village. *Eur J Gastroenterol Hepatol*. 1998;10:47-50. Crossref
12. Roussos, A., Philippou, N. and Gourgoulianis, K.I. Helicobacter pylori Infection and Respiratory Diseases: A Review. *World J Gastroenterol* 2003;9: 5-8. Crossref
13. Jia, E.Z., Zhao, F.J., Hao, B., et al. Helicobacter pylori Infection Is Associated with Decreased Serum Levels of High Density Lipoprotein, but Not with the Severity of Coronary Atherosclerosis. *Lipids in Health and Disease*, 2009;23:59. Crossref
14. Atherton, J. The Pathogenesis of H. pylori Induced Gastro-Duodenal Diseases. *Annual Review of Pathology: Mechanisms of Disease* 2006;1:63-96. Crossref
15. Price AB. The Sydney system. *Histological division. J Gastroenterol Hepatol*. 1991;6:209-22. Crossref
16. Suk F, Lien G, Yu T, Ho Y. Global trends in Helicobacter pylori research from 1991-2008 analyzed with the Science Citation Index Expanded. *Eur J Gastroenterol Hepatol*. 2011;23:295-301. Crossref
17. Sharma SK, Maharjan DK, Thapa PB. Hospital based analytic study of peptic ulcer disease in patients with dyspeptic symptoms. *Kathmandu Univ Med J* 2009;7:135-8.
18. Poudel A., Regmi S., Poudel S., Joshi P. Correlation between endoscopic and histopathological findings in gastric lesions. *Journal of Universal College of Medical Sciences* 2013;1: 37-41. Crossref

19. Ereelep OB, Caglar E, Dobrucali A. The prevalence of gastroesophageal reflux disease among hospital employees. *Dis Esophagus*. 2014;27:403-8. Crossref
20. Shrestha R, Koirala K, K C Shiva Raj, Hada Batajoo K. Helicobacter Pylori Infection Among Patients with Upper Gastrointestinal Symptoms: Prevalence and Relation to Endoscopy Diagnosis and Histopathology. *J Family Med Prim Care*. 2014;3:154–8. Crossref
21. Dixon MF. Helicobacter pylori and peptic ulceration: Histopathological aspects. *J Gastroenterol Hepatol*. 1991;6:125–30. <http://dx.doi.org/10.1111/j.1440-1746.1991.tb01451.x>
22. Zhang C, Yamada N, Wu YL, Wen M, Matsuhisa T, Matsukura N. Comparison of Helicobacter pylori infection and gastric mucosal histological features of gastric ulcer patients with chronic gastritis patients. *World J Gastroenterol*. 2005;11:976–81. Crossref
23. Chan WY, Hui PK, Leung KM. Coccoid forms of Helicobacter pylori in human stomach. *Am J Clin Pathol*. 1994;102:503–7. Crossref
24. Chen X Y, Liu WZ, Shi Y, Zhang DZ, Xiao SD, Tytgat GNJ. Helicobacter pylori associated gastric diseases and lymphoid tissue hyperplasia in gastric antral mucosa. *J Clin Pathol*. 2002;55:133–7. Crossref
25. Genta RM, Hammer HW. The significance of lymphoid follicles in the interpretation of gastric biopsy specimens. *Arch Pathol Lab Med* 1994;118:740–3. Crossref