



## Morphological and elemental analysis of termite mound and ant nest in agriculturally prominent area

Pandit Shelke<sup>1</sup>, Meghmala Waghmode<sup>1,\*</sup>, Ravindra Mene<sup>1</sup>, Aparna Gunjal<sup>2</sup>, Neha Patil<sup>1</sup>, Namdeo Bhujbal<sup>1</sup>, Urmila Dhangar<sup>1</sup>, Shital Jagtap<sup>1</sup>, Shubhangi Shinde<sup>1</sup>

<sup>1</sup>PDEA's Annasaheb Magar Mahavidyalaya, Hadapsar, Pune 411028, Maharashtra, India

<sup>2</sup>Department of Microbiology, Dr. D. Y. Patil Arts, Commerce and Science College, Pimpri, Pune 411018, Maharashtra, India

(Received: 15 August 2022; Revised: 22 March 2023; Accepted: 16 May 2023)

### Abstract

Soil management is important for the farmers to improve the crop yield. In nature some invertebrates serve as bioindicators and biomonitors. Biogenic structure built by insects is important for controlling soil erosion and water reserves. Ants and termites nest architecture along with the elemental analysis was studied to evaluate soil health and possible threats imposed by heavy metals in the area. The soil samples were collected and analyzed for various parameters. Systematic study of porosity, composition, and nutritional values of soil in ant nest and termite mound were done. The Atomic Absorption Spectrophotometer and Inductively Coupled Plasma Mass Spectrophotometer studies showed that ant nest and termite mound samples were found to contain elements viz., zinc, selenium, lead, cadmium, nickel, chromium and titanium. Based on Scanning Electron Microscope-Energy Dispersive Analysis of X-rays, the size of soil samples collected from ant nest and termite were found to be 27.77 nm and 25.56 nm, respectively. The corrosion resistant zirconium and titanium metals were detected in 0.68 and 0.39% concentration in ant nest and termite mound samples, respectively, representing the insect house as a possible source of rich metals. The ant nest and termite mound materials contain quartz, microcline, kaolinite, and clay minerals. Ant nests and termite mounds can thus be used as hydrological indicators to address the problems of soil erosion.

**Keywords:** Bioindicators, economical, heavy metal, minerals, soil, toxicity

### Introduction

Soil functioning is important considering its role in ecosystem management (Wall et al., 2012). Soil health is getting disturbed due to soil erosion and some anthropogenic activities. Termites build their mound by the combination of quartz grains with their natural secretions (humidifying agent), vegetable debris and clay minerals (plasticizer) (Echezona et al., 2012). To evaluate the soil ecosystem, bioindicators and biomonitoring functioning is very important. In nature, some species of invertebrates have been recognized as bioindicators and biomonitors of ecology. Soil ecosystems can be assessed using sentinel species as bioindicators (Amiard-Triquet et al., 2012). Based on the changes in the ecosystem affected by natural calamities (e.g., wasps' famine, soil erosion, heavy rainfall) or anthropogenic activities, bioindicator organisms change their communal behavior (Medhi et al., 2020). Reports are available on the role of termite communities in reflecting the soil conditions including macro aggregation of soil, chemical extensive richness, biodiversity, and soil hydrological functions (Duran-Bautista et al., 2020). Bioindicators include honeybees, drosophila, wasp, termites, and ants (Chowdhury et al., 2023). Role of arthropods as bioindicators is attributed to their community-based structure, nature of predator and possibility of statistical analysis (Medhi et al., 2020). Insects which have the capacity to serve as ecological biomarkers can be studied to find the ecotoxicity of that area (Amiard-Triquet et al., 2012). The insects as

bioindicators with their mechanism to combat environmental stress are represented in Table 1.

Biomonitoring is based on the finding changes in the ecosystem by using the biodiversity data of keystone species and natural inhabitants (Ma et al., 2018). The wasps are chemicaresidents of rural as well urban areas and have been reported to serve as biomonitors due to their potential to accumulate metals. *Polistes dominulus* (paper wasps) larval fecal mass are found to contain lead which indicates that wasp has good heavy metal excretion mechanism (Urbini et al., 2006).

### Ant nests

The role of ants as ecological indicator is attributed to their nest building potential using local resources (Okrutniak and Grześ, 2021; Sorvari, 2009). Ant's nest is one of the widely studied homes in context of their composition (metal accumulation), architecture (as per the environmental factors and insects' own interest), foraging behavior and ecosystem management (Fagundes et al., 2020). The way ant finds their place to build the nest or new home after the destruction or threat imposed on their nest, the journey of nest relocation, is done by scout ants which uses "one-pass" or "two pass" strategy which relies upon pheromones (Marshall et al., 2003). Scouts use Buffon's needle for the evaluation of nest size. Reports are available, which suggest ant nest size and its architecture are dependent upon the local environmental




conditions and the role of ants in their nest (Sankovitz et al., 2021). Tunnels in the ant nest indicate the competition between the social workers in ant's community. As per the pollutant concentration, ants bring about some changes in their life cycle as well as nest size. It has been

reported that the red wood ant (*Formica aquilonia*) residing in the heavy metal polluted area of copper smelter in Southwest Finland, has built small size nest and showed lower reproduction rate (Eeva et al., 2004). The attributes of the ant nest are shown in Fig. 1.

**Table 1** Insects as bioindicators with their mechanism to combat environmental stress

Examples of bioindicator insect	Stress factor	Mechanisms	References
Ground beetle <i>Parallelomorbus laevigatus</i>	Metals (As, Cd, Cr, Pb, Ni, and Hg)	Biomagnification	Conti et al. (2017)
New Zealand (NZ) cicada ( <i>Amphipsalta</i> and <i>Notopsalta</i> )	Climate change	Fast ecological radiation	James et al. (2003); Marshall et al. (2012)
Aphids ( <i>Acaudinum centaureae</i> , <i>Aphis euphorbiae</i> , <i>A. galiiscabri</i> , <i>A. helianthemii</i> , <i>A. kelimeschi</i> , <i>A. leontodontis</i> , <i>A. molluginis</i> , <i>A. pilosellae</i> , <i>A. subnitida</i> , <i>Brachycandus lucifugus</i> , <i>Nasonovia pilosellae</i> , <i>Uroleucon obscurum</i> )	Heavy metals	Species richness	Osiadacz and Halaj (2016)
Aphid ( <i>Aphis gossypii</i> )	Heavy metals (Cd, Cu, Zn, and Pb)	Bioaccumulation	Alajmi et al. (2021)
Ant ( <i>Lasius niger</i> )	Heavy metals (Cd and Zn)	Bioaccumulation	Okrutniak and Grześ (2021)
Paper wasps ( <i>Polistes dominulus</i> )	Lead	Bioaccumulation	Urbini et al. (2006)
Ant ( <i>Lasius niger</i> )	DEHP (Diethyl hexyl phthalate)	Activation of immune state	Cuviller-Hot et al. (2014)
Red wood ant ( <i>Formica aquilonia</i> )	Heavy metals (Al, Cu, Cd, Ni, Zn, As, Pb, Hg)	Lower reproduction rate	Eeva et al. (2004)




**Figure 1** Attributes of the ant nest

### Termite mound

Mound designed by termites have stereotypical architecture with structural changes as per the requirement of temperature, wind, and food. Mounds have tunnels, open chimneys, or vent holes for the exchange of gases (King et al., 2015). Termites build nests


as shelter for protection (called as “protective trail galleries”) as well as foraging tools, considering the facets of global warming induced temperature drift (Oberst et al., 2017; Oberst et al., 2020). Termite nests have been classified into three types based on their construction pattern as given in Fig. 2.



**Figure 2** Types of termite nest (Genise, 2017)

As per the reports available, mounds made by termites have outgrowth of spire which is tilted as per the sunlight and internal channels. Number of internal channels in mounds depends upon the environmental conditions like soil, temperature and wind (Fagundes et al., 2020). Termite *Macrotermes michaelseni* mounds in Northern Namibia are cone shaped extended spire and pediment (Turner, 2000). Based on the environmental challenges and number of species, termites create turrets in the mounds by which they can allot space (queen's royal chamber) for the residence as per the species and labor (Feltell et al., 2008). Reports are available where some

turrets or fungus comb when kept for the fungi; it could assist termites for the digestion of wood or old comb (Oberst et al., 2020; Rouland-Lefevre, 2000). Termites although build nests with multiple compartments, they still show communication directly through vibration (biotremology) or indirectly (stigmergy) (Calovi et al., 2019; Oberst et al., 2017). The terminate mounds can be studied for various reasons like geochemistry, sociobiology, biotite weathering into titanium and zirconium or pedogenesis mechanism (soil formation) as given in Fig. 3.



**Figure 3** Attributes of termite mound

## Materials and Methods

### Collection of the sample

Soil samples were collected from termite mound and ant nest, located in rural area of Pune city, Maharashtra, India. The method used for sampling was grab sampling. The soil samples collected were homogeneous. The collected samples were kept in clean sterile containers at 4

°C until use for analysis. The morphological and micro morphological studies were performed of ant nest and termite mound samples. Ant nest and termite mounds architectural pattern samples were evaluated in terms of height, diameter and number of turrets (Diehl et al., 2005). The sampling points for termite mound and ant nest are shown in Fig. 4 and Fig. 5 respectively.




Figure 4 Sampling point for termite mound



Figure 5 Sampling point for ant nest

### Chemical analysis of ant nest and termite mound samples

The pH and electrical conductivity (EC) parameters were measured as per the standard methods of American Public Health Association (APHA, 1998). The pH and EC were determined using pH meter and electrical conductivity meter, respectively.

### Analysis of the metals

Metal contents and micromorphology of sample was determined using SEM with EDS (Model -JSM-6360, 20kV). For the estimation of extractable metals from ant nest and termite mound, samples were digested as per the protocol of Alghanmi et al. (2015). The sample (0.5 g) was mixed with acid mixture [HCl (37%): HNO<sub>3</sub> (70%) in the ratio 3:1]. After 12 hours the sample was digested at 120 °C for 3 hours and filtered using Whatman filter paper number 1. The metal contents were analyzed using Inductively Coupled Plasma Mass Spectrophotometer (ICP MS Spectrophotometer - EXPEC 7000) and Atomic Absorption Spectrophotometer (AAS Varian Spectra, Germany).

### Determination of size of soil

The ant nest and termite mound samples were analyzed by the X-ray diffraction (XRD) techniques as per the method of Linsen et al. (2014). Advance Bruker D8 was used in this research. This instrument uses Bragg-Brentano geometry.

## Results and Discussion

Considering the ecological role of termites and ants, current study was done on finding the similarities and differences in physicochemical attributes of nest prepared by ant (*Lasius niger*) and termite (*Coptotermes formosanus*). The chemical analysis of ant nest and termite mound samples is shown in Table 2. The EC is more in termite mound sample which was found to be  $1.764 \times 10^{-3}$ . The element composition of ant nest and termite mound samples is represented in Table 3, where the oxygen content was found to be highest in ant nest and termite mound samples which was 57.66 and 57.21 % respectively. The carbon content in ant nest and termite mound samples was 21.24 and 13.27 % respectively (Table 3). Insects, specifically ants, serve as ecosystem engineers who affect physico-chemical and biological properties of the soil. Extensive study has been conducted by ecologists and biologists on the potential role of ants and termites as bioindicators and biomonitors. Research has been conducted on the design and ecological role of ant nests and termite mounds. Architectural designs and collective behavior of insects during construction of nests has been extensively studied in biomimetics like swarm robots' construction (Werfel et al., 2014). Replacement of cement with eco-friendly material like termite mound clay soil and calcium carbide wastes as binder has been tested for the production of unfired clay bricks (Akinyemi et al., 2021).

**Table 2** Morphological and chemical analysis of the ant nest and termite mound samples


Parameters	Ant nest	Termite mound
Height (m)	3.0 ± 0.01	1.0 ± 0.00
Diameter (m)	2.0 ± 0.00	2.0 ± 0.01
Number of turrets / Open channels	15.0 ± 0.00	30.0 ± 0.00
Mean diameter of turrets	0.25 ± 0.00	0.70 ± 0.01
pH	6.23	7.38
Electrical conductivity (Ohm/m)	1.112 x 10 <sup>-3</sup> ± 0.00	1.764 x 10 <sup>-3</sup> ± 0.01

The data represents average of triplicate ± standard deviation.

### XRD analysis

The X-ray diffraction patterns of the ant nest and termite mound materials are represented in Fig. 6. The ant nest and termite mound materials mainly consist of quartz, microcline, kaolinite and clay minerals. The analysis shows that the peaks referable to quartz are more prominent in both the samples while microcline and kaolinite are the dominant clay mineral, which is consistent with tropical environment (Dowouna et al.,

2012; Horiuchi et al., 2014; Liu et al., 2019). Moreover, the structural analysis of ant nest and termite mound samples reveals the polycrystalline nature. The average crystallite size (D) is calculated using Debye-Scherrer's equation from the full-width half maximum (FWHM) of all the obtained peaks for ant nest and termite mound sample and it is found that 28 and 26 nm respectively.




**Figure 6** X-ray diffraction pattern of ant nest and termite mound sample


### SEM-EDX

The SEM images of ant nest and termite mound are shown in Fig. 7 and Fig. 8, respectively at different magnification viz., x100, x500 and x1000. SEM

micrograph showed homogenous structure with micro and macropores, correlated to the structure of ant nets and termite mound.




**Figure 7** SEM image of ant nest sample




**Figure 8** SEM image of termite mound sample

The SEM-EDAX images of ant nest and termite mound samples showed presence of precious metals viz., oxygen, carbon, copper, magnesium, aluminium, iron, zirconium,

calcium, potassium, and silica (Fig. 9 and 10). The metals platinum and titanium were also found in termite mound samples (Fig. 10).



**Figure 9** SEM-EDAX image of ant nest sample




**Figure 10** SEM-EDAX image of termite mound sample

**Table 3** Elemental composition in ant nest and termite mound samples as per SEM-EDS data


Elements	Ant nest sample	Termite mound sample
	Weight (%)	
Carbon	21.24 ± 0.01	13.27 ± 0.01
Oxygen	57.66 ± 0.00	57.21 ± 0.00
Magnesium	0.60 ± 0.02	1.69 ± 0.00
Aluminium	4.05 ± 0.01	4.11 ± 0.01
Silica	8.81 ± 0.02	11.29 ± 0.00
Potassium	0.42 ± 0.00	-
Calcium	1.68 ± 0.02	3.25 ± 0.01
Iron	4.24 ± 0.00	6.95 ± 0.00
Copper	0.63 ± 0.00	0.63 ± 0.00
Zirconium	0.68 ± 0.00	-
Zinc	-	0.90 ± 0.00
Titanium	-	0.39 ± 0.01

The data represents average of triplicate ± standard deviation.

Ants are common in metal polluted areas due to their metal resistant behavior (Klimek et al., 2022). Reports are available on the rare earth elements and trace elements in termite mound. In one study, ten termite mounds and adjacent topsoil from Central and North-Eastern Namibia was analyzed for the rare earth and mobile element. Compared to adjacent soil, *Macrotermes* termite mound has been reported to have more concentration of titanium, zirconium, selenium and Th/U ratios suggesting mineralization and chemical weathering ability of termites (Sako et al., 2009). Study conducted on ants living in meadow and forest sites in a mining and smelting region near Olkusz, Poland concluded that diversity increases with increase in metal pollution (Grześ, 2009). The metal contents (Zn, Cd, Ni, Cr, Se, and Pb) were analyzed using ICP-MS and AAS. Negligible differences were observed in the concentration of metals estimated by using AAS and ICP-MS (Figs. 11 and 12).



**Figure 11** Metal content in samples using AAS




**Figure 12** Metal content in samples using ICP-MS

### FTIR analysis of the ant nest and termite mound

The FTIR spectra of the ant nest and termite mound samples are shown in Fig. 13. The peak is observed at

1002.32 nm wavelength indicating the presence of presence of alkyl/aryl halides like F/Cl/I.



**Figure 13** FTIR spectra of the ant nest and termite mound sample

### Conclusions

In nature, different types of biogenic structures are observed, among which ant nest and termite mound have been studied in the biomimetic field. High mechanical resistance and ventilation pattern are two important aspects which are considered by ant and termite mound, while building their home. Nest architecture and its metallic as well as mineral contents were studied in this work. Both ant and termite nests, residing in close proximity, are found to have very much similarity in metal and mineral content, the exception was titanium and zirconium. This type of study is important for the metallurgical data on such biogenic structure. In drier areas of India such termite mound can be considered as hydrological indicators and will also prevent soil erosion. As compared to the nearby agricultural soil, ant nests and mounds showed more metallic contents.

**Author Contributions:** PS: Data analysis, MW: Experiments conducted and paper writing, RM: Data analysis for XRD, AG: Experiment design and paper writing, NP: paper writing, NB: FTIR analysis, UD: sample collection, SJ: sample collection, SS: SEM data analysis.

**Conflicts of Interest:** The author declares no conflicts of interest.

**Data Availability Statement:** The data that support the finding of this study are available from the corresponding author, upon reasonable request.

### References

- Akinyemi, B.A., Orogade, B.O., & Okoro, C.W. (2021). The potential of calcium carbide waste and termite mound soil as materials in the production of unfired clay bricks. *Journal of Cleaner Production*, 279, 123693.
- Alajmi, R., Al-Shammari, M., Abdel-Gaber, R., Metwally, D., El-Khadragy, M.F., & Alrajeh, A. (2021). Effectiveness of naturally occurring *Aphis gossypii* on tomato plants as a bio-indicator for heavy metals in Riyadh and Hafar Al-Batin, Saudi Arabia. *Saudi Journal of Biological Sciences*, 28(7), 4096-4101.
- Alghanmi, S.I., Al Sulami, A.F., El-Zayat, T.A., Alhogbi, B.G., & Salam, M.A. (2015). Acid leaching of heavy metals from contaminated soil collected from Jeddah, Saudi Arabia: Kinetic and thermodynamics studies. *International Soil and Water Conservation Research*, 3(3), 196-208.
- Amiard-Triquet, C., Amiard, J.C., & Rainbow, P.S. (Eds.). (2012). *Ecological biomarkers: Indicators of ecotoxicological effects*. CRC Press.
- APHA. (1998). *Standard methods for analysis of water and waste water*. The American Public Health Association, United States.
- Calovi, D.S., Bardunias, P., Carey, N., Scott Turner, J., Nagpal, R., & Werfel, J. (2019). Surface curvature guides early construction activity in mound-building termites. *Philosophical Transactions of the Royal Society B*, 374(1774), 20180374.
- Chowdhury, S., Dubey, K., Choudhury, S., Das, A., Jeengar, D., Sujatha, B., Kumar, A., Kumar, N.,



- Semwal, A., & Kumar, V. (2023). Insects as bioindicator: A hidden gem for environmental monitoring. *Frontiers in Environmental Science*, 11, 1146052. doi 10.3389/fenvs.2023.1146052.
- Conti, E., Dattilo, S., Costa, G., & Puglisi, C. (2017). The ground beetle *Parallelomorpha laevigatus* is a potential indicator of trace metal contamination on the eastern coast of Sicily. *Ecotoxicology and Environmental Safety*, 135, 183-190.
- Cuviller-Hot V., Salin K., Devers, S., Tasiemski, A., Schaffner, P., Boulay, R., Biliard, S., & Lenoir, A. (2014). Impact of ecological doses of the most widespread phthalate on a terrestrial species, the ant *Lasius niger*. *Environmental Research*, 131, 104-110.
- Diehl, E., Junqueira, L.K., & Berti-Filho, E. (2005). Ant and termite mound co-inhabitants in the wetlands of Santo Antonio da Patrulha, Rio Grande do Sul, Brazil. *Brazilian Journal of Biology*, 65(3), 431-437.
- Dowouna, G., Pearl, A., Dubbin, W., Prosper, N., Baba, M., Eric, N., & Richard, H. (2012). Characteristics of termite mounds and associated Acrisols in the coastal savanna zone of Ghana and impact on hydraulic conductivity. *Natural Science*, 4, 423-437. doi 10.4236/ns.2012.47058.
- Duran-Bautista, E.H., Armbrecht, I., Acioli, A.N.S., Suárez, J.C., Romero, M., Quintero, M., & Lavelle, P. (2020). Termites as indicators of soil ecosystem services in transformed amazon landscapes. *Ecological Indicators*, 117, 106550.
- Echezona, B.C., Igwe, C.A., & Attama, L.A. (2012). Properties of arboreal ant and ground-termite nests in relation to their nesting sites and location in a tropical-derived Savanna. *Psyche: A Journal of Entomology*, 235840. doi 10.1155/2012/235840.
- Eeva, T., Sorvari, J., & Koivunen, V. (2004). Effects of heavy metal pollution on red wood ant (*Formica* s. str.) populations. *Environmental Pollution*, 132(3), 533-539.
- Fagundes, T.M., Ordóñez, J.C., & Yaghoobian, N. (2020). How the thermal environment shapes the structure of termite mounds. *Royal Society Open Science*, 7, 191332. doi 10.1098/rsos.191332.
- Feltell, D., Bai, L., & Jensen, H.J. (2008). An individual approach to modelling emergent structure in termite swarm systems. *International Journal of Modelling, Identification and Control*, 3(1), 29-40.
- Genise, J.F. (2017). Blueprints of termite and ant nests. In J.F. Genise (Ed.), *Ichnoentomology: Insect Traces in Soils and Paleosols*. Springer, pp. 247-284. doi 10.1007/978-3-319-28210-7\_11.
- Grześ, I.M. (2009). Ant species richness and evenness increase along a metal pollution gradient in the Boleslaw zinc smelter area. *Pedobiologia*, 53(1), 65-73.
- Horiuchi, Y., Ohno, T., Hoshino, M., Shin, K.-C., Murakami, H., Tsunematsu, M., & Watanabe, Y. (2014). Geochemical prospecting for rare earth elements using termite mound materials. *Mineralium Deposita*, 49, 1013-1023. doi 10.1007/s00126-014-0550-3.
- James, M., Kovacs, T., Anna, D., & Nigel, F. (2003). Simulating the evolution of ant behavior in evaluating nest sites. In: Banzhaf, W., Ziegler, J., Christaller, T., Dittrich, P., & Kim, J.T. (eds.) *Advances in Artificial Life. Lecture Notes in Computer Science*, Vol 2801. Springer, Berlin, Heidelberg. doi 10.1007/978-3-540-39432-7\_69.
- King, H., Ocko, S., & Mahadevan, L. (2015). Termite mounds harness diurnal temperature oscillations for ventilation. *PNAS*, 112(37), 11589-11593.
- Klimek, B., Poliwka-Modliborek, H., & Grześ, I.M. (2022). Ant nests as a microbial hot spots in a long-term heavy metal-contaminated soils. *Environmental Science and Pollution Research*, 29, 10848-10857. doi 10.1007/s11356-021-16384-y.
- Linsen, D.G., Xuefa, S.I., Yanguang, L.U., Xisheng, F.G., Zhihua, C.N., Chunjuan, W.G., Jianjun, Z.U., & Yuanhui, H.G. (2014). Mineralogical study of surface sediments in the western Arctic Ocean and their implications for material sources. *Polar Research*, 25(3), 192-203.
- Liu, X., Zhang, C., Guang, Y., & Yanfeng, G. (2019). Bioinspired ant-nest-like hierarchical porous material using CaCl<sub>2</sub> as additive for smart indoor humidity control. *Industrial and Engineering Chemistry Research*, 58(17), 7139-7145. doi 10.1021/acs.iecr.8b06092.
- Ma, A., Bohan, D.A., Canard, E., Derocles, S.A., Gray, C., Lu, X., & Kratina, P. (2018). A replicated network approach to 'big data' in ecology. In Bohan, D.A., Dumbbell, A.J., Woodward, G., & Jackson, M. (Eds.) *Advances in Ecological Research* (Vol. 59, pp. 225-264). Academic Press.
- Marshall, D.C., Hill, K.B., Marske, K.A., Chambers, C., Buckley, T.R., & Simon, C. (2012). Limited, episodic diversification and contrasting phylogeography in a New Zealand cicada radiation. *BMC Evolutionary Biology*, 12(1), 1-18.
- Medhi, J., Dutta, J., & Kalita, M.C. (2020). Biomonitoring ecosystem: modeling relationship with arthropods. In Ranz, R.E.R. (Ed.), *Arthropods- Are they beneficial for mankind?* IntechOpen. doi 10.5772/intechopen.94313.
- Oberst, S., Bann, G., Lai, J.C., & Evans, T.A. (2017). Cryptic termites avoid predatory ants by eavesdropping on vibrational cues from their footsteps. *Ecology Letters*, 20(2), 212-221.
- Oberst, S., Lai, J.C., Martin, R., Halkon, B.J., Saadatfar, M., & Evans, T.A. (2020). Revisiting stigmergy in light of multi-functional, biogenic, termite structures as communication channel. *Computational and Structural Biotechnology Journal*, 18, 2522-2534.
- Okrutniak, M., & Grześ, I.M. (2021). Accumulation of metals in *Lasius niger*: Implications for using ants as bioindicators. *Environmental Pollution*, 268, 115824.

- Osiadacz, B., & Halaj, R. (2016). Aphids under stress. Species groups and ecological functional groups of aphids define heavy metal grasslands of Central Europe. *Redia, XCLA*, 35-51.
- Rouland-Lefèvre, C. (2000). Symbiosis with fungi. In Abe, T., Bignell, D.E. & Higashi, M. (Eds.), *Termites: Evolution, sociality, symbioses, ecology*. Kluwer Academic, Dordrecht, The Netherlands, pp. 289-306.
- Sako, A., Mills, A.J., & Roychoudhury, A.N. (2009). Rare earth and trace element geochemistry of termite mounds in central and northeastern Namibia: Mechanisms for micro-nutrient accumulation. *Geoderma*, 153(1-2), 217-230.
- Sankovitz, M., & Purcell, J. (2021). Ant nest architecture is shaped by local adaptation and plastic response to temperature. *Scientific Reports*, 11, 23053. doi 10.1038/s41598-021-02491-w.
- Sorvari, J. (2009). Foraging distances and potentiality in forest pest insect control: An example 480 with two candidate ants (*Hymenoptera*: Formicidae). *Myrmecological News*, 12, 211-215.
- Turner, J.S. (2000). Architecture and morphogenesis in the mound of *Macrotermes michaelseni* (Sjostedt) (*Isoptera*: Termitidae, Macrotermitinae) in Northern Namibia. *Cimbebasia*, 16, 143-175.
- Urbini, A., Sparvoli, E., & Turillazzi, S. (2006). Social paper wasps as bioindicators: A preliminary research with *Polistes dominulus* (*Hymenoptera vespidae*) as a trace metal accumulator. *Chemosphere*, 64(5), 697-703.
- Wall, D.H., Behan-Pelletier, V., Jones, T.H., Ritz, K., Six, J., Strong, D.R., & van der Putten, W.H. (2012). Soil ecology and ecosystem services. Oxford University Press, Oxford Academic. doi 10.1093/acprof:oso/9780199575923.001.0001.
- Werfel, J., Petersen, K., & Nagpal, R. (2014). Designing collective behavior in a termite-inspired robot construction team. *Science*, 343, 754-758.